

Appendix B

Adult Prevalence, Initiation, and Cessation

	<u>Page</u>
Maryland	82
Allegany	84
Anne Arundel	86
Baltimore City	88
Baltimore County	90
Calvert	92
Caroline	94
Carroll	96
Cecil	98
Charles	100
Dorchester	102
Frederick	104
Garrett	106
Harford	108
Howard	110
Kent	112
Montgomery	114
Prince George's	116
Queen Anne's	118
St. Mary's	120
Somerset	122
Talbot	124
Washington	126
Wicomico	128
Worcester	130



Maryland

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	21.5% ± 0.9% 822,418	19.6% ± 1.0% 760,986	<u>17.7%</u> ± 0.8% 711,036	*16.1% ± 0.8% 655,763
Any Tobacco Product (Adult Minority Populations)	19.7% ± 1.8% 256,851	18.3% ± 1.8% 260,298	15.8% ± 1.5% 239,905	17.4% ± 1.6% 247,871
Cigarette	16.9% ± 0.9% 648,628	<u>14.8%</u> ± 0.9% 580,696	13.8% ± 0.8% 556,955	*12.4% ± 0.7% 510,440
Cigar	5.9% ± 0.5% 225,452	5.9% ± 0.6% 233,399	5.1% ± 0.5% 207,383	*4.0% ± 0.5% 163,755
Smokeless Tobacco	1.1% ± 0.2% 41,541	1.0% ± 0.2% 40,874	1.1% ± 0.2% 45,234	1.3% ± 0.2% 53,455

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	6.2% ± 2.9% 10,727	7.6% ± 3.3% 11,350	9.5% ± 4.5% 15,267	7.0% ± 3.8% 8,557

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	23.4% ± 3.4% 73,606	22.2% ± 3.7% 67,716	<u>31.7%</u> ± 4.1% 88,244	29.7% ± 3.5% 95,037

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Selected Populations

Proportion (%) of Selected Populations and Number For Pregnant Women


Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Pregnant Women 100% OF Maryland Birth Certificates	9.2% 6,842	8.0% 5,877	6.8% 5,261	6.6% 5,105
Female	15.2%	13.1%	11.8%	10.7%
Male	18.8%	16.8%	15.9%	14.3%
Black/African American	19.1%	15.5%	14.5%	14.4%
Hispanic	18.3%	16.1%	9.8%	11.7%
White	16.6%	14.4%	14.1%	11.6%

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Proportion with Adult Smokers	33.9% ± 1.8% 511,692	31.9% ± 1.8% 521,430	29.3% ± 1.6% 493,787	*23.9% ± 1.4% 424,458

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Allegany County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	22.6% ± 4.6% 16,733	23.9% ± 4.5% 14,097	22.2% ± 4.6% 11,769	26.8% ± 5.0% 13,884
Any Tobacco Product (Adult Minority Populations)	16.6% ± 11.9% 2,083	29.0% ± 17.7% 1,714	39.3% ± 21.4% 2,106	NOT AVAILABLE—TOO FEW MINORITY RESPONDENTS
Cigarette	19.1% ± 4.3% 14,147	19.5% ± 4.2% 11,521	18.2% ± 4.4% 9,627	20.4% ± 4.6% 10,698
Cigar	6.0% ± 2.7% 4,443	6.0% ± 2.8% 3,554	5.1% ± 2.3% 2,717	4.2% ± 3.6% 2,213
Smokeless Tobacco	2.8% ± 1.9% 2,071	4.5% ± 2.1% 2,677	4.2% ± 2.7% 2,207	7.4% ± 4.1% 3,873

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	1.4% ± 2.1% 65	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	34.1% ± 23.7% 1,872	26.9% ± 16.5% 1,651	37.8% ± 23.8% 2,065	22.5% ± 13.4% 1,405

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Pregnant Women Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Allegany County	16.8% 133	18.1% 128	22.1% 157	22.7% 153	24.8% 159	20.2% 131	23.7% 166	19.0% 130	24.7% 176

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	37.9% ± 10.9% 10,495	40.3% ± 9.2% 7,434	37.8% ± 9.4% 7,158	44.6% ± 9.0% 8,278

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Anne Arundel County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	23.5% ± 2.9% 86,281	19.7% ± 3.0% 71,322	19.1% ± 2.5% 69,560	*14.9% ± 2.3% 54,839
Any Tobacco Product (Adult Minority Populations)	23.4% ± 7.5% 15,028	20.7% ± 7.9% 15,422	16.8% ± 5.7% 11,867	16.4% ± 5.9% 12,490
Cigarette	18.6% ± 2.7% 68,495	13.5% ± 2.6% 49,284	14.0% ± 2.2% 51,344	11.5% ± 2.0% 42,329
Cigar	7.1% ± 1.8% 26,308	7.3% ± 2.0% 26,656	7.1% ± 1.6% 25,928	4.0% ± 1.4% 14,696
Smokeless Tobacco	1.5% ± 0.9% 5,396	1.0% ± 0.7% 3,797	1.3% ± 0.7% 4,865	1.7% ± 0.8% 6,378

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	19.1% ± 16.4% 3,317	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	28.0% ± 10.2% 9,137	18.1% ± 9.9% 4,810	28.9% ± 11.6% 7,071	24.9% ± 10.4% 6,592

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Anne Arundel County	11.1% 754	10.5% 724	9.7% 656	8.5% 589	8.7% 587	7.6% 514	8.1% 576	7.7% 547	7.7% 552

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	37.3% ± 5.3% 54,805	31.8% ± 5.4% 47,214	27.8% ± 4.5% 42,400	23.2% ± 4.2% 36,921

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Baltimore City

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	32.5% ± 3.8% 130,207	27.3% ± 3.4% 131,971	22.3% ± 2.9% 99,574	*20.8% ± 2.8% 93,888
Any Tobacco Product (Adult Minority Populations)	33.9% ± 5.1% 86,508	28.3% ± 4.5% 90,644	24.1% ± 3.9% 71,855	*23.6% ± 3.7% 71,644
Cigarette	28.6% ± 3.7% 114,784	23.8% ± 3.3% 115,871	19.5% ± 2.8% 87,645	*16.7% ± 2.5% 75,678
Cigar	5.4% ± 1.6% 21,709	6.2% ± 2.0% 30,157	4.0% ± 1.4% 17,868	5.0% ± 1.7% 22,683
Smokeless Tobacco	0.6% ± 0.6% 2,594	0.6% ± 0.5% 2,811	NOT AVAILABLE—TOO FEW SMOKLESS USING RESPONDENTS	0.4% ± 0.4% 1,720

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	11.1% ± 10.5% 1,831	7.1% ± 8.0% 1,639	9.9% ± 11.7% 1,938	13.2% ± 13.0% 2,127

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	14.3% ± 6.7% 7,074	11.3% ± 5.9% 6,470	25.8% ± 9.0% 13,286	21.7% ± 8.6% 11,254

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Baltimore City	14.9% 1,435	14.6% 1,328	13.9% 1,254	13.4% 1,214	12.0% 1,106	11.6% 1,064	10.3% 1,009	9.9% 997	10.3% 1,025

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	48.1% ± 6.9% 70,800	49.8% ± 6.5% 89,794	43.7% ± 6.2% 72,072	*32.7% ± 5.5% 58,066

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Baltimore County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	21.9% ± 2.7% 133,311	21.8% ± 2.9% 124,628	20.7% ± 3.1% 99,574	*16.4% ± 2.4% 96,027
Any Tobacco Product (Adult Minority Populations)	17.9% ± 4.6% 32,985	20.9% ± 6.1% 29,849	18.4% ± 4.6% 30,361	17.5% ± 4.6% 31,720
Cigarette	17.2% ± 2.4% 105,142	15.8% ± 2.6% 90,914	17.5% ± 2.9% 101,876	13.7% ± 2.3% 80,722
Cigar	6.0% ± 1.5% 36,791	6.8% ± 1.7% 38,909	5.4% ± 1.9% 31,767	*3.2% ± 1.2% 18,869
Smokeless Tobacco	0.7% ± 0.5% 4,257	0.8% ± 0.6% 4,595	1.1% ± 1.0% 6,606	0.9% ± 0.8% 5,458

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	5.6% ± 6.2% 1,620	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	11.1% ± 12.8% 3,942	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	33.0% ± 9.8% 19,762	23.2% ± 10.5% 12,102	36.1% ± 13.7% 16,443	41.4% ± 11.4% 19,153

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Baltimore County	9.6% 904	8.9% 807	8.8% 791	8.8% 822	8.6% 809	7.8% 758	7.8% 776	7.6% 769	7.3% 748

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	35.0% ± 5.4% 72,542	35.1% ± 5.4% 79,680	32.5% ± 6.2% 66,740	26.2% ± 4.8% 60,729

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Calvert County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	25.8% ± 4.0% 13,091	26.3% ± 4.4% 13,670	21.6% ± 4.5% 13,747	20.7% ± 4.2% 13,352
Any Tobacco Product (Adult Minority Populations)	30.4% ± 10.8% 2,628	15.3% ± 9.3% 1,520	21.1% ± 11.5% 1,979	20.8% ± 13.8% 1,523
Cigarette	20.0% ± 3.6% 10,178	17.8% ± 3.8% 9,260	17.2% ± 4.3% 11,020	16.0% ± 3.9% 10,481
Cigar	7.1% ± 2.4% 3,622	10.8% ± 3.3% 5,651	5.7% ± 2.6% 3,666	4.4% ± 2.0% 2,860
Smokeless Tobacco	1.5% ± 1.1% 779	1.1% ± 0.8% 574	1.4% ± 1.1% 885	1.4% ± 1.2% 910

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	12.9% ± 12.6% 372	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	22.3% ± 11.6% 1,154	10.4% ± 10.1% 442	20.6% ± 13.9% 1,209	25.8% ± 13.2% 1,851

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Calvert County	14.2% 145	14.3% 137	12.7% 129	12.2% 129	11.8% 118	11.9% 119	12.7% 128	9.9% 97	11.5% 110

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	38.9% ± 6.7% 10,387	41.8% ± 7.2% 10,673	30.5% ± 7.3% 8,805	30.7% ± 6.7% 10,497

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Caroline County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	26.7% ± 4.1% 5,883	23.1% ± 4.6% 4,976	19.4% ± 5.1% 4,530	25.8% ± 7.5% 6,228
Any Tobacco Product (Adult Minority Populations)	24.2% ± 11.5% 807	19.2% ± 10.3% 1,005	28.5% ± 16.4% 1,333	41.2% ± 26.6% 1,429
Cigarette	22.6% ± 3.8% 5,029	18.3% ± 4.0% 3,952	15.5% ± 4.5% 3,585	20.3% ± 7.1% 4,922
Cigar	6.4% ± 2.5% 1,423	5.4% ± 2.4% 1,181	6.8% ± 3.1% 9,392	7.4% ± 6.8% 1,797
Smokeless Tobacco	1.8% ± 1.2% 405	2.7% ± 1.7% 583	0.5% ± 0.6% 116	2.2% ± 1.8% 526

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	24.5% ± 12.9% 461	17.5% ± 12.2% 265	NOT AVAILABLE—TOO FEW QUITTING RESPONDENTS	16.1% ± 14.7% 476

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Caroline County	17.0% 69	18.3% 71	15.1% 58	16.2% 68	15.9% 74	17.8% 76	12.8% 61	13.7% 64	12.8% 64

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	43.2% \pm 7.1% 4,237	34.9% \pm 8.9% 3,103	31.9% \pm 9.4% 3,109	32.3% \pm 8.4% 3,433

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Carroll County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	26.2% ± 4.5% 26,432	19.1% ± 4.0% 20,761	20.3% ± 3.6% 24,756	14.4% ± 3.0% 17,698
Any Tobacco Product (Adult Minority Populations)	NOT AVAILABLE—TOO FEW TOBACCO USING RESPONDENTS	NOT AVAILABLE—TOO FEW TOBACCO USING RESPONDENTS	28.3% ± 15.6% 1,924	17.0% ± 16.8% 1,569
Cigarette	18.3% ± 4.0% 18,616	11.4% ± 3.0% 12,349	13.8% ± 3.0% 16,844	10.5% ± 2.6% 12,956
Cigar	9.6% ± 2.9% 9,763	8.5% ± 3.0% 9,223	7.7% ± 2.8% 9,392	4.2% ± 1.9% 5,209
Smokeless Tobacco	2.3% ± 1.8% 2,355	1.6% ± 1.3% 1,713	2.2% ± 1.7% 2,725	2.3% ± 1.4% 2,795

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	24.7% ± 14.6% 2,092	50.0% ± 21.1% 3,776	29.6% ± 18.4% 2,438	36.2% ± 16.0% 2,768

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Carroll County	11.1% 211	10.6% 199	11.2% 213	9.6% 188	9.6% 191	9.6% 184	10.2% 191	11.1% 207	10.6% 186

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	43.2% ± 7.1% 4,237	34.9% ± 8.9% 3,103	31.9% ± 9.4% 3,109	32.3% ± 8.4% 3,433

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Cecil County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	27.1% ± 4.7% 16,792	29.5% ± 4.7% 17,980	25.3% ± 3.8% 17,990	23.6% ± 4.3% 17,342
Any Tobacco Product (Adult Minority Populations)	33.4% ± 18.6% 2,645	47.5% ± 18.0% 2,746	30.2% ± 17.5% 1,980	9.8% ± 6.5% 675
Cigarette	23.0% ± 4.5% 14,274	22.9% ± 4.3% 14,045	21.3% ± 3.6% 15,320	19.3% ± 4.0% 14,325
Cigar	7.0% ± 2.9% 4,373	6.8% ± 2.8% 4,234	7.0% ± 2.7% 5,038	6.2% ± 2.9% 4,565
Smokeless Tobacco	2.8% ± 1.6% 1,750	1.4% ± 1.1% 853	1.7% ± 1.1% 1,234	2.0% ± 1.3% 1,512

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	13.1% ± 10.5% 927	35.8% ± 14.9% 2,524	23.6% ± 10.3% 1,576	28.2% ± 12.8% 2,536

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Cecil County	23.3% 265	19.5% 230	18.2% 211	18.9% 230	19.7% 239	18.7% 227	19.7% 267	21.4% 283	22.8% 291

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	40.5% ± 8.2% 10,313	45.1% ± 7.6% 13,240	39.7% ± 6.3% 12,514	37.2% ± 7.4% 12,995

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Charles County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	24.6% ± 4.4% 19,539	23.0% ± 4.2% 19,712	17.5% ± 3.3% 17,372	21.1% ± 3.5% 21,326
Any Tobacco Product (Adult Minority Populations)	17.8% ± 7.3% 4,087	17.3% ± 6.5% 5,020	13.4% ± 5.3% 5,208	18.6% ± 5.7% 6,536
Cigarette	18.9% ± 3.9% 15,111	17.9% ± 3.9% 15,401	12.9% ± 2.9% 12,840	16.1% ± 3.1% 16,396
Cigar	7.0% ± 2.8% 5,657	5.6% ± 2.2% 4,797	4.9% ± 1.8% 4,882	7.0% ± 2.4% 7,059
Smokeless Tobacco	0.7% ± 0.7% 535	1.8% ± 1.4% 1,519	1.8% ± 1.4% 1,815	1.9% ± 1.4% 1,978

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	12.3% ± 11.1% 901	17.3% ± 13.5% 1,121	38.8% ± 15.0% 2,854	25.4% ± 11.2% 2,432

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Charles County	13.3% 232	12.2% 213	11.1% 195	9.4% 177	10.7% 194	7.1% 132	8.7% 169	6.8% 134	7.0% 134

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	35.6% ± 7.9% 13,362	35.6% ± 7.2% 14,684	31.3% ± 6.1% 14,898	30.5% ± 5.6% 15,450

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Dorchester County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	27.4% ± 6.2% 6,728	23.5% ± 4.7% 5,501	19.0% ± 4.9% 4,482	19.5% ± 4.2% 4,748
Any Tobacco Product (Adult Minority Populations)	23.6% ± 8.5% 1,335	13.4% ± 7.9% 923	19.7% ± 12.2% 1,243	23.6% ± 9.8% 1,421
Cigarette	24.9% ± 6.2% 6,173	19.8% ± 4.4% 4,643	16.4% ± 4.7% 3,907	15.3% ± 3.7% 3,736
Cigar	5.2% ± 2.3% 1,295	4.2% ± 2.1% 978	3.0% ± 1.8% 723	3.2% ± 1.7% 779
Smokeless Tobacco	0.4% ± 0.5% 92	1.6% ± 1.6% 384	1.3% ± 1.1% 315	2.1% ± 1.7% 504

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	20.3% ± 13.6% 669	12.8% ± 9.7% 271	<u>47.8%</u> ± 22.0% 983	27.6% ± 15.4% 801

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Dorchester County	18.6% 61	14.3% 48	16.6% 51	14.9% 52	13.2% 48	17.5% 57	13.1% 53	11.4% 47	14.4% 65

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	35.6% ± 7.9% 13,362	35.6% ± 7.2% 14,684	31.3% ± 6.1% 14,898	30.5% ± 5.6% 15,450

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Frederick County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	23.6% ± 3.8% 34,152	19.2% ± 4.0% 26,852	18.7% ± 3.1% 28,967	17.0% ± 3.2% 27,008
Any Tobacco Product (Adult Minority Populations)	27.5% ± 13.3% 4,905	13.8% ± 9.3% 2,520	12.8% ± 7.6% 2,285	19.4% ± 10.5% 3,854
Cigarette	16.3% ± 3.3% 23,680	13.7% ± 3.6% 19,212	11.6% ± 2.5% 18,149	12.2% ± 2.8% 19,562
Cigar	8.1% ± 2.5% 11,820	6.9% ± 2.5% 9,762	7.3% ± 2.1% 11,493	3.9% ± 1.8% 6,158
Smokeless Tobacco	2.8% ± 1.6% 4,075	1.7% ± 1.2% 2,434	2.4% ± 1.4% 3,821	2.1% ± 1.3% 3,323

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	19.2% ± 12.6% 2,326	24.7% ± 15.5% 2,660	25.7% ± 14.3% 1,908	24.5% ± 12.8% 3,007

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Frederick County	12.1% 351	12.0% 349	9.2% 276	9.6% 297	8.6% 254	8.0% 241	7.5% 231	7.3% 224	7.4% 219

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	35.0% ± 7.0% 20,570	32.3% ± 7.1% 20,506	29.2% ± 5.6% 21,057	23.1% ± 5.2% 18,514

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Garrett County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	25.2% ± 5.0% 6,088	22.7% ± 4.4% 5,065	22.7% ± 4.3% 5,078	15.3% ± 4.0% 3,424
Any Tobacco Product (Adult Minority Populations)	NOT AVAILABLE—TOO FEW MINORITY RESPONDENTS	NOT AVAILABLE—TOO FEW MINORITY RESPONDENTS	NOT AVAILABLE—TOO FEW MINORITY RESPONDENTS	NOT AVAILABLE—TOO FEW MINORITY RESPONDENTS
Cigarette	19.3% ± 4.9% 4,678	16.4% ± 3.9% 3,653	16.5% ± 3.8% 3,702	10.8% ± 3.6% 2,411
Cigar	6.8% ± 2.7% 1,657	5.6% ± 2.3% 1,260	4.4% ± 2.1% 981	2.4% ± 2.1% 532
Smokeless Tobacco	6.5% ± 2.7% 1,573	4.3% ± 2.2% 965	5.9% ± 2.5% 1,317	5.3% ± 2.5% 1,189

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	14.6% ± 12.1% 237	24.6% ± 15.7% 408	30.6% ± 21.6% 375	44.4% ± 21.8% 769

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Garrett County	16.8% 56	19.2% 63	17.0% 51	16.7% 57	20.9% 66	19.4% 61	16.2% 48	18.1% 57	17.0% 47

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	23.3% ± 9.2% 2,250	36.5% ± 8.6% 2,955	30.8% ± 8.4% 2,709	21.2% ± 7.8% 1,809

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Harford County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	22.2% ± 4.1% 34,284	21.2% ± 4.1% 32,854	22.0% ± 3.0% 38,323	<u>15.1%</u> ± 3.4% 26,841
Any Tobacco Product (Adult Minority Populations)	22.4% ± 12.0% 4,927	20.2% ± 11.5% 3,967	21.5% ± 7.9% 5,379	17.9% ± 9.8% 4,318
Cigarette	16.7% ± 3.7% 25,951	14.8% ± 3.5% 23,184	17.4% ± 2.8% 30,444	<u>11.4%</u> ± 2.9% 20,163
Cigar	8.9% ± 3.1% 13,849	7.1% ± 2.7% 11,158	6.5% ± 2.0% 11,408	4.4% ± 2.2% 7,797
Smokeless Tobacco	1.5% ± 1.2% 2,303	1.4% ± 1.2% 2,239	2.0% ± 1.3% 3,565	0.9% ± 0.7% 1,584

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	11.5% ± 13.6% 1,250	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	22.2% ± 15.9% 2,249	16.9% ± 15.2% 1,715	25.2% ± 10.3% 3,649	44.1% ± 19.6% 6,251

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Harford County	13.1% 386	10.1% 291	12.4% 360	11.6% 337	10.1% 301	10.7% 314	11.3% 342	10.8% 327	9.1% 271

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	36.9% ± 7.3% 24,219	30.2% ± 6.8% 22,244	31.9% ± 5.1% 24,769	*20.6% ± 5.4% 16,855

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Howard County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	16.2% ± 3.0% 31,691	15.3% ± 3.0% 26,901	12.5% ± 2.4% 23,867	*8.3% ± 1.7% 16,518
Any Tobacco Product (Adult Minority Populations)	17.3% ± 7.0% 8,335	12.8% ± 5.7% 5,584	10.0% ± 4.8% 5,857	*4.6% ± 2.2% 2,937
Cigarette	11.3% ± 2.7% 22,158	9.5% ± 2.6% 16,711	7.9% ± 2.1% 15,264	*5.8% ± 1.5% 11,660
Cigar	6.2% ± 1.9% 12,154	7.1% ± 2.4% 12,608	6.0% ± 1.8% 11,568	*2.6% ± 1.0% 5,145
Smokeless Tobacco	0.9% ± 0.6% 1,799	1.9% ± 1.8% 3,376	0.2% ± 0.2% 409	0.6% ± 0.4% 1,171

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	22.5% ± 12.3% 2,181	29.6% ± 15.6% 2,457	30.6% ± 15.3% 2,042	38.1% ± 15.0% 3,853

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Howard County	3.3% 119	4.4% 158	3.3% 117	3.6% 128	3.3% 116	3.7% 124	2.9% 99	2.7% 93	2.2% 74

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	26.7% ± 5.3% 23,104	27.1% ± 5.4% 23,244	19.9% ± 4.4% 18,602	*15.3% ± 3.5% 14,026

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Kent County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	22.5% ± 5.5% 2,845	20.9% ± 5.3% 3,122	18.8% ± 5.8% 2,716	21.0% ± 5.9% 3,037
Any Tobacco Product (Adult Minority Populations)	20.7% ± 12.2% 405	22.5% ± 16.9% 706	29.8% ± 21.7% 777	13.8% ± 11.1% 244
Cigarette	18.3% ± 5.3% 2,335	16.2% ± 5.1% 2,432	15.1% ± 5.5% 2,200	16.1% ± 4.9% 2,339
Cigar	4.3% ± 2.2% 549	6.7% ± 2.7% 1,014	3.6% ± 2.2% 529	3.7% ± 2.2% 547
Smokeless Tobacco	4.5% ± 4.6% 582	0.6% ± 0.7% 97	2.2% ± 2.1% 323	5.3% ± 4.3% 772

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	32.2% ± 20.1% 437	23.4% ± 17.2% 356	NOT AVAILABLE—TOO FEW QUITTING RESPONDENTS	35.0% ± 25.4% 598

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Kent County	21.2% 43	20.0% 31	16.6% 26	18.6% 31	16.3% 33	15.5% 28	19.0% 36	19.7% 35	13.7% 30

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	41.0% ± 9.2% 1,784	38.1% ± 9.9% 1,962	38.9% ± 11.8% 2,233	33.6% ± 11.7% 1,696

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Montgomery County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	14.0% ± 2.4% 92,026	14.2% ± 2.7% 90,464	10.8% ± 2.1% 72,961	10.1% ± 2.0% 69,638
Any Tobacco Product (Adult Minority Populations)	11.6% ± 4.1% 27,151	14.6% ± 5.5% 33,167	9.4% ± 3.6% 26,067	12.8% ± 4.4% 25,477
Cigarette	8.8% ± 1.9% 58,227	9.9% ± 2.4% 64,088	7.5% ± 1.8% 51,176	7.2% ± 1.6% 49,788
Cigar	4.9% ± 1.5% 32,338	5.2% ± 1.7% 33,825	4.0% ± 1.4% 27,328	*2.2% ± 1.1% 14,900
Smokeless Tobacco	0.5% ± 0.4% 3,241	0.4% ± 0.5% 2,567	0.5% ± 0.5% 3,409	0.6% ± 0.5% 4,449

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	25.2% ± 13.0% 8,174	31.5% ± 17.3% 11,947	35.2% ± 16.5% 10,928	34.3% ± 13.6% 12,073

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Montgomery County	2.5% 327	2.0% 269	1.3% 168	1.3% 171	1.0% 142	0.9% 120	0.7% 95	0.5% 68	0.6% 80

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	19.9% ± 4.4% 51,360	19.4% ± 4.7% 52,278	20.8% ± 4.5% 59,233	14.0% ± 3.4% 43,342

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Prince George's County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	16.2% ± 2.9% 82,026	12.9% ± 2.4% 74,437	13.9% ± 2.4% 82,441	16.4% ± 2.5% 97,354
Any Tobacco Product (Adult Minority Populations)	14.2% ± 3.4% 48,842	11.2% ± 2.8% 48,760	12.5% ± 2.7% 58,441	16.2% ± 3.0% 70,666
Cigarette	13.9% ± 2.7% 70,493	10.7% ± 2.3% 62,385	10.8% ± 2.1% 64,438	12.6% ± 2.2% 75,780
Cigar	3.7% ± 1.5% 18,621	3.3% ± 1.2% 19,301	3.6% ± 1.4% 21,789	4.9% ± 1.6% 29,255
Smokeless Tobacco	0.1% ± 0.2% 731	0.5% ± 0.5% 2,669	0.8% ± 0.6% 5,086	1.0% ± 0.7% 5,847

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	12.8% ± 13.5% 2,653	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	22.3% ± 12.3% 9,045	25.0% ± 11.7% 8,599	34.0% ± 13.8% 10,646	24.6% ± 10.4% 10,801

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Prince George's County	3.6% 447	3.1% 386	2.5% 316	2.1% 269	1.9% 237	1.6% 205	1.4% 177	1.4% 185	1.7% 213

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	32.6% ± 5.7% 70,153	25.5% ± 5.7% 66,752	26.2% ± 4.5% 71,933	23.8% ± 4.5% 61,378

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Queen Anne's County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	26.8% ± 4.7% 8,897	<u>17.7%</u> ± 3.8% 5,338	21.5% ± 4.9% 7,387	20.8% ± 5.3% 7,322
Any Tobacco Product (Adult Minority Populations)	22.2% ± 12.5% 1,027	16.7% ± 13.7% 604	21.5% ± 16.0% 1,354	26.9% ± 23.4% 902
Cigarette	23.6% ± 4.5% 7,838	<u>14.3%</u> ± 3.5% 4,318	15.4% ± 4.2% 5,303	14.6% ± 4.7% 5,100
Cigar	6.1% ± 3.2% 2,019	3.3% ± 1.6% 1,011	<u>9.5%</u> ± 3.9% 3,266	6.1% ± 3.5% 2,142
Smokeless Tobacco	1.8% ± 1.4% 607	1.9% ± 1.4% 587	3.9% ± 2.9% 1,340	4.3% ± 2.5% 1,523

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	17.4% ± 10.4% 568	34.7% ± 16.9% 803	49.4% ± 28.9% 1,730	40.3% ± 17.4% 1,588

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Queen Anne's County	15.6% 78	11.7% 55	10.2% 54	8.0% 40	10.8% 55	9.8% 51	8.9% 46	11.3% 63	6.2% 33

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	40.2% ± 7.9% 5,328	35.1% ± 7.5% 4,143	28.1% ± 8.6% 3,766	37.6% ± 9.1% 5,933

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





St. Mary's County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	26.5% ± 4.2% 17,031	19.4% ± 3.8% 11,865	18.9% ± 3.9% 12,659	20.0% ± 3.7% 14,002
Any Tobacco Product (Adult Minority Populations)	22.7% ± 11.9% 2,428	10.4% ± 6.9% 1,305	11.7% ± 7.1% 1,216	19.8% ± 10.4% 1,764
Cigarette	20.0% ± 3.8% 12,862	14.8% ± 3.4% 9,166	15.0% ± 3.6% 10,106	15.6% ± 3.5% 11,035
Cigar	8.4% ± 2.6% 5,409	4.9% ± 2.0% 2,986	4.2% ± 1.8% 2,830	4.1% ± 1.9% 2,895
Smokeless Tobacco	2.6% ± 1.6% 1,690	1.0% ± 0.9% 622	2.1% ± 1.4% 1,415	2.0% ± 1.3% 1,415

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	21.1% ± 11.3% 1,311	26.6% ± 17.0% 1,230	54.3% ± 17.2% 2,945	34.5% ± 17.5% 1,767

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
St. Mary's County	13.0% 158	14.3% 181	12.3% 165	11.8% 159	13.5% 195	11.5% 160	9.9% 148	10.5% 155	8.8% 127

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	37.8% ± 6.9% 11,082	38.2% ± 7.4% 10,628	29.6% ± 7.1% 9,208	22.7% ± 5.6% 7,463

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Somerset County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	25.0% ± 4.5% 5,207	20.2% ± 4.4% 4,058	24.7% ± 7.8% 4,260	27.2% ± 7.3% 4,568
Any Tobacco Product (Adult Minority Populations)	22.6% ± 7.9% 1,538	19.8% ± 7.1% 1,794	21.1% ± 15.8% 971	26.5% ± 12.4% 886
Cigarette	19.8% ± 4.0% 4,128	18.1% ± 4.2% 3,633	19.9% ± 7.3% 3,450	22.6% ± 7.4% 3,857
Cigar	4.9% ± 2.2% 1,014	4.1% ± 2.2% 827	4.5% ± 2.8% 783	7.9% ± 6.5% 1,315
Smokeless Tobacco	2.0% ± 1.5% 416	0.9% ± 0.9% 188	3.4% ± 3.2% 591	1.7% ± 1.8% 289

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	28.5% ± 13.8% 566	24.2% ± 14.1% 373	33.2% ± 30.2% 586	25.0% ± 18.7% 580

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Somerset County	20.9% 57	23.0% 56	17.0% 44	19.7% 51	16.5% 44	16.2% 43	16.1% 45	14.8% 37	14.8% 41

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	44.8% ± 9.0% 3,226	37.4% ± 9.8% 2,268	46.6% ± 17.0% 3,055	37.3% ± 10.1% 2,268

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Talbot County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	16.5% ± 5.5% 4,716	25.0% ± 6.7% 6,467	19.3% ± 5.1% 5,298	21.1% ± 6.1% 5,984
Any Tobacco Product (Adult Minority Populations)	10.1% ± 10.3% 769	40.8% ± 21.0% 2,430	18.0% ± 13.6% 595	7.9% ± 7.2% 289
Cigarette	13.8% ± 4.9% 3,983	18.2% ± 5.8% 4,780	13.5% ± 4.2% 3,742	17.9% ± 5.8% 5,087
Cigar	4.4% ± 2.4% 1,286	5.6% ± 2.2% 1,480	6.2% ± 3.4% 1,752	4.0% ± 3.4% 1,142
Smokeless Tobacco	0.3% ± 0.5% 93	2.6% ± 3.5% 675	0.6% ± 0.6% 163	1.9% ± 2.4% 534

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	17.6% ± 12.8% 358	14.3% ± 11.8% 389	45.4% ± 23.3% 1,086	NOT AVAILABLE—TOO FEW QUITTING RESPONDENTS

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Talbot County	14.4% 53	10.1% 36	10.2% 34	10.3% 35	8.0% 31	12.9% 47	10.1% 36	8.2% 30	10.6% 41

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	31.1% ± 9.5% 2,476	41.3% ± 10.7% 3,210	31.8% ± 11.7% 3,107	29.7% ± 11.5% 2,980

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Washington County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	27.1% ± 4.7% 25,607	24.6% ± 4.5% 24,602	19.1% ± 3.1% 19,053	*18.8% ± 3.5% 18,864
Any Tobacco Product (Adult Minority Populations)	35.7% ± 18.7% 3,633	31.6% ± 17.4% 4,330	16.6% ± 9.1% 1,042	21.0% ± 14.1% 1,772
Cigarette	21.7% ± 4.4% 20,799	18.8% ± 3.9% 18,936	15.8% ± 2.9% 15,848	*13.0% ± 3.1% 13,227
Cigar	5.7% ± 2.6% 5,397	7.3% ± 2.8% 7,356	4.3% ± 1.6% 4,264	4.4% ± 1.9% 4,416
Smokeless Tobacco	2.5% ± 1.9% 2,424	3.4% ± 2.1% 3,465	1.0% ± 0.7% 1,030	3.3% ± 1.8% 3,327

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW QUITTING RESPONDENTS	16.5% ± 15.2% 1,268	29.7% ± 13.6% 1,980	31.8% ± 16.3% 2,201

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Washington County	19.1% 305	20.9% 339	18.1% 306	17.5% 296	15.6% 266	15.8% 278	16.3% 310	16.3% 322	16.3% 298

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	40.5% ± 8.7% 15,369	31.2% ± 7.9% 12,602	28.7% ± 5.9% 12,548	*25.3% ± 6.0% 11,071

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Wicomico County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	24.3% ± 5.1% 13,999	22.6% ± 4.3% 14,055	24.1% ± 5.2% 15,589	19.4% ± 4.1% 13,070
Any Tobacco Product (Adult Minority Populations)	18.1% ± 7.5% 2,541	22.1% ± 9.6% 3,627	22.3% ± 11.2% 3,818	22.1% ± 10.4% 3,678
Cigarette	20.9% ± 5.0% 12,093	19.7% ± 4.1% 12,361	18.7% ± 4.6% 12,170	16.7% ± 3.9% 11,262
Cigar	4.0% ± 2.0% 2,290	4.2% ± 1.9% 2,622	5.8% ± 3.1% 3,762	6.2% ± 2.9% 4,154
Smokeless Tobacco	1.8% ± 1.6% 1,062	2.1% ± 1.7% 1,310	1.9% ± 1.8% 1,216	2.0% ± 1.9% 1,360

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	3.0% ± 3.8% 152	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	28.2% ± 14.6% 1,520	25.7% ± 14.0% 1,721	7.7% ± 7.8% 241	23.1% ± 15.7% 1,468

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates



Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Wicomico County	14.4% 169	16.6% 189	15.1% 175	15.7% 186	14.8% 172	16.7% 202	13.3% 177	15.2% 204	15.5% 212

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	34.9% \pm 9.4% 8,085	34.2% \pm 8.1% 8,296	31.5% \pm 8.6% 8,238	26.7% \pm 7.0% 8,710

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)





Worcester County

Tobacco Use by Adults: 2000—2008

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Any Tobacco Product (All Adult Populations)	24.8% ± 5.9% 8,852	28.0% ± 4.8% 10,290	22.3% ± 5.1% 8,518	22.8% ± 6.6% 8,799
Any Tobacco Product (Adult Minority Populations)	9.8% ± 9.2% 761	25.6% ± 12.9% 1,918	32.1% ± 16.9% 1,973	<u>7.5%</u> ± 6.0% 338
Cigarette	20.7% ± 5.3% 7,452	23.3% ± 4.6% 8,596	18.0% ± 4.6% 6,955	17.7% ± 6.0% 6,926
Cigar	4.6% ± 2.2% 1,667	7.7% ± 2.8% 2,848	5.3% ± 3.0% 2,042	6.7% ± 4.5% 2,628
Smokeless Tobacco	2.0% ± 1.5% 710	0.5% ± 0.6% 175	2.0% ± 2.0% 782	2.6% ± 2.9% 1,018

INITIATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE—TOO FEW INITIATING RESPONDENTS

CESSATION OF TOBACCO USE IN PAST YEAR

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
Adult Population	13.9% ± 11.3% 339	10.4% ± 8.6% 361	36.4% ± 20.6% 1,158	NOT AVAILABLE—TOO FEW QUITTING RESPONDENTS

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2008 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women

Proportion (%) of Selected Populations and Number For Pregnant Women from 100% of Birth Certificates

Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008
Maryland Statewide	9.2%	8.7%	8.0%	7.7%	7.4%	6.9%	6.8%	6.6%	6.6%
Worcester County	16.9% 84	13.8% 72	15.4% 70	16.4% 77	14.9% 67	12.1% 60	16.1% 75	14.0% 70	14.2% 68

ADULT HOUSEHOLDS WITH MINOR CHILDREN

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008
<i>Proportion with Adult Smokers</i>	34.3% ± 16.6% 4,627	44.8% ± 10.0% 4,366	37.5% ± 11.1% 4,974	42.9% ± 13.1% 5,445

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)

