



STATE OF MARYLAND
DHMH

Maryland Department of Health and Mental Hygiene
201 W. Preston Street • Baltimore, Maryland 21201

Martin O'Malley, Governor – Anthony G. Brown, Lt. Governor – Joshua M. Sharfstein, M.D., Secretary

NOV 16 2011

The Honorable Martin O'Malley
Governor
State of Maryland
Annapolis, MD 21401-1991

The Honorable Thomas V. Mike Miller, Jr.
President of the Senate
H-107 State House
Annapolis, MD 21401-1991

The Honorable Michael E. Busch
Speaker of the House
H-101 State House
Annapolis, MD 21401-1991

Re: Health-General Article, §13-1004(d), FY 2011 Biennial Tobacco Study
Cigarette Restitution Fund – Tobacco Use Prevention and Cessation Program

Dear Governor O'Malley, President Miller, and Speaker Busch:

Pursuant to Health-General Article, §13-1004(d), Annotated Code of Maryland, the Department of Health and Mental Hygiene is directed to produce a biennial legislative report on the results of the Biennial Tobacco Study.

The enclosed legislative report summarizes the findings from the Fiscal Year 2011 Maryland Youth Tobacco Survey (MYTS) and data from the 2000-2010 Maryland Behavioral Risk Factor Surveillance System Survey (BRFSS), as available. This report also uses Fiscal Year 2009 data from the Maryland Adult Tobacco Survey (MATS). The MYTS and MATS surveys were administered in each county and Baltimore City, resulting in estimates of tobacco use by underage middle school and high school youth as well as for adults age 18 and older in each of those jurisdictions and Statewide. The significant findings from the surveys document continued reductions in tobacco use behaviors since program inception in Fiscal Year 2001.

Thank you for your commitment to the progress we are making in reducing tobacco use in Maryland. If you have questions about this report, please contact Ms. Marie Grant, Director of Governmental Affairs at (410) 767-6481.

Sincerely,

Joshua M. Sharfstein, M.D.
Secretary

Enclosure

cc: Ms. Sarah Albert, MSAR #2866
Marie Grant, J.D.
Frances Phillips, R.N., M.H.A.

Donna Gugel, M.H.S.
Donald Shell, M.D., M.S.



Monitoring Changing Tobacco Use Behaviors 2000—2010

Legislative Report

Maryland Department of Health & Mental Hygiene

**Cigarette Restitution Fund's
Tobacco Use Prevention and Cessation Program**

Fiscal Year 2011

Martin O'Malley
Governor
State of Maryland

Anthony G. Brown
Lieutenant Governor

Joshua M. Sharfstein, MD
Secretary
Department of Health & Mental Hygiene

Frances B. Phillips, RN, MHA
Deputy Secretary, Public Health Services
Department of Health & Mental Hygiene

Donna Gugel, MHS
Interim Director, Family Health Administration
Department of Health & Mental Hygiene

Donald Shell, MD, MA
Interim Director, Center for Health Promotion & Education
Family Health Administration, Department of Health & Mental Hygiene



Table of Contents

	<u>Pages</u>
Highlights.....	1
Introduction.....	4
Health and Tobacco.....	6
Underage Youth.....	8
Adults.....	18
Secondhand Smoke.....	30
References.....	32
Appendices.....	33
Appendix A: Youth Prevalence, Initiation & Cessation	34
Appendix B: Adult Prevalence, Initiation & Cessation	86

Sources




Unless otherwise explicitly stated, the source for all data pertaining to youth comes from the Maryland Youth Tobacco Surveys (2000, 2002, 2006, 2008, and 2010). With respect to adults, unless otherwise explicitly stated, the source for all data pertaining to adults comes from the Behavioral Risk Factor Surveillance Survey (2000 –2010) or the Maryland Adult Tobacco Surveys (MATS) (2000, 2002, 2006, and 2008). All data for pregnant women comes from the Maryland Office of Vital Statistics.

Statutory Authority and Requirements


Maryland’s Health-General Article, Title 13, Subtitle 10, requires that the Maryland Department of Health and Mental Hygiene conduct a biennial tobacco study and report specific findings to the Maryland General Assembly and the Governor. The appendices to this Executive Summary provide the detailed data for indicators on which DHMH is required by Subtitle 10 to report as part of its biennial tobacco study. Indicators pertaining to youth and drawn from youth surveys are set forth in Appendix A. Indicators pertaining to adults and drawn from adult surveys are set forth in Appendix B.

Between 2000 and 2010 Maryland made substantial progress in reducing the use of cigarettes by underage youth, adults, and pregnant women.


- ☛ 38.9% decrease in cigarette use among all youth (from 15.7% in 2000 to 9.6% in 2010).
- ☛ 25.9% decrease in cigarette use among adults (from 20.5% in 2000 to 15.2% in 2010).
- ☛ 33.7% decrease in smoking among pregnant women (from 9.2% in 2000 to 6.1% in 2010).
- ☛ Cigarettes and cigars continue to be the most popular tobacco products among underage high school current tobacco users, 17.5% of whom use both products. In addition, 15.9% of underage high school current tobacco users use cigarettes, cigars, as well as other tobacco products (bidis, kreteks, pipes, or smokeless tobacco).




Maryland’s Clean Indoor Air Act prohibits smoking indoors at work and in restaurants and bars and has a 94% compliance rate. Voluntary indoor smoking bans at home and in cars are benefiting underage Maryland youth.



- ☞ 16.3% increase in the proportion of underage youth living in smoking households that prohibit smoking inside the household (from 48.5% in 2006 to 56.4% in 2010).
- ☞ 11.1% increase in the proportion of underage youth living in non-smoking households that prohibit smoking inside the household (from 84.0% in 2006 to 93.3% in 2010).



- ☞ 65.9% increase in underage youth not exposed to secondhand smoke indoors (from 44.3% in 2000 to 73.5% in 2010).



- ☞ 28.8% increase in underage youth not exposed to secondhand smoke while riding in a vehicle (from 57.6% in 2000 to 74.2% in 2010).

- ☞ Between 2000 and 2010, adult cigarette smoking in Maryland decreased significantly. Maryland has the 11th lowest rate of adult cigarette smoking in the nation.
- ☞ Because of differences in surveys used by states, it is not possible to rank states' underage youth cigarette smoking as can be done for adults. Between 2000 and 2010, cigarette smoking decreased by 38.7% among underage Maryland high school youth and by 51.4% among underage Maryland middle school youth.

Changes in Use of Selected Tobacco Products Maryland: 2000-2010

Cigarette Smoking:  **51.4% decrease among underage middle school youth (from 7.2% in 2000 to 3.5% in 2010)**

 **38.7% decrease among underage high school youth (from 23.0% in 2000 to 14.1% in 2010)**

 **25.9% decrease among adults (from 20.5% in 2000 to 15.2% in 2010)**

 **33.7% decrease among pregnant women (from 9.2% in 2000 to 6.1% in 2010)**

Cigar Smoking:  **23.9% decrease among underage middle school youth (from 4.6% in 2000 to 3.5% in 2010)**

 **11.2% increase among underage high school youth (from 12.5% in 2000 to 13.9% in 2010)**

Smokeless

Tobacco Use:  **38.1% decrease among underage middle school youth (from 2.1% in 2000 to 1.3% in 2010)**

 **2.1% increase among underage high school youth (from 4.7% in 2000 to 4.8% in 2010)**

In Fiscal Year 2001, using non-tax proceeds from participation in the national Master Settlement Agreement with the tobacco industry, Maryland implemented a comprehensive tobacco control program modeled on the best practice recommendations of the Centers for Disease Control and Prevention. In addition, Maryland increased its excise tax on cigarettes to \$2.00 per pack and implemented a comprehensive statutory smoking ban that applies to all worksites, restaurants, and bars. Due to the combination of programmatic and policy initiatives undertaken since 1992, Maryland has the 11th lowest adult prevalence of cigarette smoking in the nation.

This document constitutes the report to the Maryland General Assembly and the Governor on the FY 2011 Tobacco Study mandated by Health—General Article Title 13, Subtitle 10. The appendices to this report include Statewide and county-specific data on changing tobacco use behaviors, including those measures required of the FY 2011 Tobacco Study report. Appendix A focuses on underage youth behaviors whereas Appendix B focuses on adult behaviors.

ABOUT THE DATA

The data presented in this report use findings from: (1) the Maryland Youth Tobacco Survey (MYTS); (2) the Maryland Behavioral Risk Factor Surveillance System Survey (BRFSS); and (3) the Maryland Adult Tobacco Survey (MATS). In previous reports, the adult data source was exclusively MATS. This report, however, uses comparable adult BRFSS data from 2000-2010, as available.

The following important differences between MATS and BRFSS data exist, and should be kept in mind when reviewing this report:

- ◆ BRFSS was designed to provide Statewide estimates of relevant behaviors. BRFSS data are weighted so that data collected reflect the population of the State as a whole, not the county where the data are collected. BRFSS county-level estimates should be used with this fact in mind.
- ◆ MATS was designed to provide both county-specific as well as Statewide estimates of tobacco use behaviors. MATS survey data were weighted so that data collected in a county reflected the population of that county. When all the county data were aggregated, they then matched the Statewide population.


- ◆ MATS collects data during the last quarter of every other calendar year. BRFSS collects data throughout each calendar year. MATS data provide a snapshot of tobacco use behaviors during the last quarter of the year conducted, while BRFSS data represent an average of those behaviors across an entire calendar year.
- ◆ There are substantial differences in survey size. MATS normally had in excess of 22,000 adult respondents per year. At present BRFSS typically includes 8,800 respondents, and during the first half of the decade, included on average 4,400 respondents per year. The smaller sample size of BRFSS makes data analysis of sub-populations and small and medium-sized jurisdictions challenging because of the smaller number of survey respondents from those jurisdictions.

Part 1 — Health and Tobacco


In Maryland, cigarette smoking is the risk behavior that causes the greatest number of preventable deaths every year (6,800),¹ with almost 150,000 additional adults suffering daily from one or more diseases caused by smoking.² These numbers do not include those who become sick or die as a result of smoking cigars, using other tobacco products, or from exposure to secondhand smoke.

It is clear that the implementation of proven evidence-based prevention programs and policies reduces tobacco-related cancers and disease, and eliminates expenditures that would be necessary to treat those diseases.

The annual cost of treating cancers and disease caused by cigarette smoking alone in Maryland was estimated to be \$2.2 billion in 2004, and will likely increase to between \$3.5-\$5.0 billion annually by 2015.¹ The human and economic toll that smoking and tobacco use exacts from Maryland residents, the Maryland economy, and the Maryland budget, can be reduced through a long-term sustained effort to reduce the use of tobacco products in this State. The cost of treating disease caused by cigarette smoking inflates the tax burden on the average Maryland household by an estimated \$622 every year.




The impact of smoking and tobacco use on health is not limited to only those persons who have been diagnosed with a cancer or disease caused by smoking, it also affects the general health status of both adults and youth. Figure 2 below shows that for both youth and adults, a higher proportion of those who have never smoked consistently report better current health status as compared to current smokers based on the 2010 survey.



Part 2 — Underage Youth


Every year, almost 90% of Marylanders who start using tobacco for the first time are youth under 18 years old. For that reason, ‘Part 2’ of this report focuses on changes in the tobacco use behaviors of underage youth. All references to “youth” in this Part refer only to underage youth (under 18 years old). The figure below presents a simplified logic model of how tobacco control policies and Maryland’s comprehensive tobacco use prevention and cessation program work together to reduce the initiation and use of tobacco products among underage Maryland youth.



Current Use of Tobacco Products


The prevalence of tobacco use is significantly higher among high school youth than middle school youth (23.8% and 7.9%, respectively). The Centers for Disease Control and Prevention (CDC) defines the “current” use of any tobacco product by youth as any use during the prior 30 days. The following figures focus primarily on tobacco use behaviors of high school youth. Detailed data on middle school tobacco use behaviors can be found in Appendix A.

In 2010, cigarettes were used by 14.1% of Maryland high school youth. Cigars and smokeless tobacco products were used by 13.9% and 4.8% of these youth, respectively. Decreases in cigarette and cigar smoking between 2008 and 2010 were statistically significant.



Concurrent Use of Tobacco Products

In 2010, 23.8% of underage high school youth used some form of tobacco and the majority of these youth used cigarettes (60.3%) or cigars (57.2%). In addition, 43.5% of underage high school tobacco users used tobacco products other than cigarettes or cigars (predominately smokeless tobacco, but also bidis, kreteks, or pipes). Figure 4 presents the percentage of underage high school tobacco users who use different types of tobacco products.



Of the 60.3% of Underage High School Current Tobacco Users Who Smoke Cigarettes:

- A. 21.0% smoke cigarettes only.
- B. 17.5% smoke both cigarettes and cigars, but do not use other tobacco products.
- D. 5.9% smoke cigarettes and use other tobacco products, but do not smoke cigars.
- C. 15.9% smoke cigarettes, cigars, and use other tobacco products.

Of the 57.2% of Underage High School Current Tobacco Users Who Smoke Cigars:

- E. 18.0% smoke cigars only.
- B. 17.5% smoke both cigars and cigarettes, but do not use other tobacco products.
- F. 5.8% smoke cigars and use other tobacco products, but do not smoke cigarettes.
- C. 15.9% smoke cigars, cigarettes, and use other tobacco products.


Of the 43.5% of Underage High School Current Tobacco Users Who Use Other Tobacco Products (Bidis, Kreteks, Pipes, or Smokeless Tobacco):

- G. 15.9% use other tobacco products only – other than cigarettes and cigars.
- D. 5.9% use other tobacco products and smoke cigarettes, but do not smoke cigars.
- F. 5.8% use other tobacco products and smoke cigars, but do not smoke cigarettes.
- C. 15.9% smoke cigarettes, cigars, and use other tobacco products.

Although cigarettes are still the most commonly used tobacco product, many young tobacco users use them concurrently with cigars or other tobacco products. Additionally, 39.7% of underage Maryland high school tobacco users do not use cigarettes. This shows the importance of prevention programs addressing the use of all tobacco products and not focusing exclusively on cigarettes.

Current Use of Flavored Tobacco


In 2010, 13.9% of underage high school youth reported smoking at least one cigar during the preceding 30 days, of whom a substantial majority (76.2%) reported that they had smoked a flavored cigar. In addition, of the 4.8% of underage high school youth who reported using smokeless tobacco during the preceding 30 days, over two-thirds (69.3%) reported that they had used flavored smokeless tobacco.



Underage Access to Tobacco Products

In Maryland as throughout the nation, it is illegal to sell cigarettes or other tobacco products to youth under 18 years old. The 2010 Regulations Restricting the Sale and Distribution of Cigarettes and Smokeless Tobacco to Protect Children and Adolescents require retailers to check photo identification of anyone purchasing cigarettes or smokeless tobacco who is under the age of 27. Frequent cigarette smokers (defined by the CDC as youth reporting that they smoked cigarettes on between 20 and 30 days during the past month) are the most likely to use retail cigarette outlets to acquire cigarettes.



In 2010, 4.4% of underage high school youth reported smoking cigarettes frequently (i.e., on 20 or more of the past 30 days) and 61.9% of these youth attempted to purchase cigarettes from a retail outlet. This represents a statistically significant decline from the 68.3% who reported purchase attempts in 2008. Among the youth attempting to purchase cigarettes from retail outlets, only 33.6% reported that they were asked for identification to prove that they were old enough to purchase cigarettes. Of the youth who were not asked for identification at a retail outlet, 74.8% were able to purchase cigarettes despite their age. In contrast, of the youth who were asked for identification, significantly fewer (34.2%) were sold cigarettes.



Underage Smoking and Alcohol and Marijuana Use

Figures 7 and 8 show that underage cigarette smoking is highly correlated with both underage drinking and the use of marijuana. Underage high school youth who smoke cigarettes are 10 times more likely to drink alcohol and 15 times more likely to smoke marijuana than are non-smokers.


Approximately 80% of underage high school smokers report they currently use alcohol (i.e., have had one or more drinks of alcohol in the past 30 days) as compared to just 25% of non-smokers. Similarly, 70% of smokers reported they currently use marijuana (i.e., have smoked marijuana during the past 30 days) as compared to just 13.3% of non-smokers.



Current alcohol and marijuana use is also highly correlated with cigarette smoking. 57.7% of smokers reported using both alcohol and marijuana during the previous 30 days as compared to 8.5% of non-smokers.

Concurrent Use of Cigarettes, Alcohol, and Marijuana

In 2010, 35.5% of underage high school youth used cigarettes, alcohol, or marijuana. Figure 9 presents the percentage of these youth who exclusively used any of these three products or a combination thereof.



Of the Underage High School Youth Who Report Currently Using Cigarettes, Alcohol, and/or Marijuana:

- A. 4.0% smoke cigarettes only.
- E. 37.3% drink alcohol only.
- G. 9.7% use marijuana only.
- B. 6.3% smoke cigarettes and drink alcohol, but do not use marijuana.
- D. 3.6% smoke cigarettes and marijuana, but do not drink alcohol.
- F. 19.3% drink alcohol and use marijuana, but do not smoke cigarettes.
- C. 19.8% smoke cigarettes, drink alcohol, and use marijuana.


Underage Youth Exposure to Tobacco-Use Prevention Messages in the Media

The tobacco industry spends an estimated \$144 million a year in Maryland to advertise and promote its products according to the latest estimates from the Campaign for Tobacco Free Kids.³ With respect to underage youth and tobacco advertising, Congress found that children are exposed to substantial tobacco advertising and that they are more influenced by tobacco marketing than are adults.

While the CDC continues to recommend that health messaging to counter tobacco industry advertising be a part of every state's tobacco control program, national, state, and local media-based youth prevention initiatives continue to decrease. Consequently, fewer Maryland high school youth are exposed to tobacco-use prevention messages. Between 2000 and 2010, the percentage of underage high school youth exposed to tobacco prevention messages in the media declined significantly by 21%.


In 2009, Congress enacted the Family Smoking Prevention and Tobacco Control Act, a federal law that gives the U.S. Food and Drug Administration the power to regulate the tobacco industry. This federal act recognizes that the tobacco industry's marketing and promotion practices are more influential on tobacco choices of youth than those of adults. Congressional findings made in connection with the act included:

- ◆ Children are exposed to substantial tobacco advertising, leading to positive beliefs about tobacco use, playing a role in leading them to overestimate the prevalence of tobacco use, and increasing the number who begin to use tobacco; and
- ◆ Children are more influenced by tobacco marketing than adults. More than 80% of youth smoke three heavily marketed brands, while only 54% of adults age 26 and older smoke these brands.





School-Based Tobacco Use Prevention Curricula

In 2010, 74.7% of underage Maryland public middle school youth reported that they had been taught about the dangers of tobacco use during the preceding school year (2009-2010). In contrast, 52.0% of underage Maryland public high school youth reported having been taught about the dangers of tobacco use during the preceding school year. Between 2008 and 2010, however, there was a significant increase in the percentage of high school youth who reported receiving this type of instruction.




Underage Youth Attitudes & Beliefs

School-based health prevention education, community-based prevention activities, restrictions on youth access to tobacco, Statewide prevention campaigns, mass media prevention messages, and tobacco industry marketing, promotion, and advertising all influence the attitudes of underage youth towards smoking and tobacco products. Between 2008 and 2010, there were significant declines in the percentage of underage high school youth who believe that smoking helps youth to “fit in or look cool,” among both non-smokers and smokers. There was no significant change, however, in the percentage of underage high school youth who believe that smokers have more friends, regardless of smoking status.



Part 3 — Adult Tobacco Use

One of the primary goals of Maryland’s comprehensive tobacco use prevention and cessation program is to reduce tobacco use by Maryland adults by motivating tobacco-users to quit and by providing assistance to those who try to quit successfully. The figure below presents a simplified logic model of how tobacco control policies and Maryland’s comprehensive tobacco use prevention program work together to reduce the use of tobacco products through cessation.



Current Use of Tobacco Products

The Centers for Disease Control and Prevention (CDC) define a current cigarette smoker as an adult who reports having smoked at least 100 cigarettes (5 packs) in their lifetime and also reports having smoked a cigarette during the past 30 days. Maryland has made significant progress in reducing adult cigarette smoking. Overall, current cigarette use among adults declined significantly by 26% between 2000 and 2010. However, smoking prevalence has remained essentially unchanged since 2008.

Changes in Cigarette Smoking from 2000—2010

Changes in Cigarette Smoking within Maryland Adult Populations:

- ⬇️ 25.9% Among all adults (from 20.5% in 2000 to 15.2% in 2010)*
- ⬇️ 54.7% Among Hispanic adults (from 17.2% in 2000 to 7.8% in 2010) ^
- ⬇️ 28.6% Among non-Hispanic White adults (from 21.0% in 2000 to 15.0% in 2010)*
- ⬇️ 33.7% Among pregnant women (from 9.2% in 2000 to 6.1% in 2010)
- ⬇️ 18.0% Among non-Hispanic Black adults (from 21.7% in 2000 to 17.8% in 2010)^


Source: Behavioral Risk Factor Surveillance Survey, 2000— 2010 and Maryland Vital Statistics Administration Birth Statistics, 2000-2010.

* Statistically significant result

^ Statistically non-significant result

Note: Cigarette smoking rates for pregnant women are based on actual birth certificate records rather than sample survey data so statistical significance does not apply.


Figure 14
Current Cigarette Use
 Maryland Adults, 2000—2010





Source: Behavioral Risk Factor Surveillance Surveys, 2000—


Like many health risk behaviors, cigarette use varies by socio-demographic characteristics and, nationwide, Black or African-American adults exhibit higher incidences of lung cancer than any other racial/ethnic group. Between 2000 and 2010, current cigarette use among Black or African-American Maryland adults declined by a statistically non-significant 18%. Comparing current cigarette use among Maryland Hispanic adults for the 2000—2004 and 2006—2010 periods reveals that there was a significant decline of 40.3% between the two five-year periods. In addition, between 2000 and 2010, there was a significant decline of 29% in current cigarette use among White adults in Maryland.

In the figures that follow, the 90% confidence interval width is presented as a line within each bar to enable the reader to assess whether apparent increases or decreases are statistically significant. If the lines representing the confidence interval of the years being compared overlap, then the difference between those years is not statistically significant. Larger confidence intervals reflect smaller sample sizes as compared to populations with smaller confidence intervals. Due to small sample sizes of Maryland Hispanic adults in each year of the Behavioral Risk Factor Surveillance Survey, Figure 16 presents combined smoking prevalence estimates for two five-year periods, 2000—2004 and 2006—2010.








Tobacco use is inversely associated with educational attainment. Maryland adults who did not graduate from high school are approximately five times more likely to be current smokers than adults who graduated from college.





An examination of rates of current cigarette use by race and educational attainment reveals that within each level of educational attainment, there are no significant differences between Black or African-American and White adults.




Between 2000 and 2010, current cigarette use among Maryland adults with a college degree declined significantly by 39%, whereas the rates of current cigarette use among adults who did not graduate from high school, have only a high school diploma, or attended some college did not exhibit any significant change.






Between 2000 and 2010, among adults who have ever smoked cigarettes, there has been a significant decline in the percentage who are current smokers. Consequently, there has been a significant increase in the percentage of former smokers.



There has also been a modest yet statistically significant increase in the percentage of Maryland adults who have never smoked cigarettes, from 56.0% in 2000 to 60.9% in 2010.




Reasons for Wanting to Quit Smoking

This section of the report focuses on adult smoking cessation. It presents data from the 2008 Maryland Adult Tobacco Survey, the latest data available on smoking cessation by adults in Maryland.

The reasons for trying to quit smoking are varied and have changed in their relative importance over time. In 2008, current and former smokers were asked to identify all their reasons for quitting (former smokers) or trying to quit (current smokers). Each respondent was allowed to identify multiple reasons for trying to quit. Next, respondents were asked to select the single most important reason to them for wanting to quit cigarette smoking.

<u>Top 5 Most Cited Reasons for Quitting</u>	<u>2000</u>	<u>2008</u>	<u>% Change</u>
When identifying multiple reasons for quitting			
1. The cost of tobacco products	31.0%	48.9%	+58%
2. Physical fitness	35.7%	48.2%	+35%
3. Health hazards associated with smoking	29.3%	46.9%	+60%
4. Encouragement from a friend or relative	23.4%	39.5%	+69%
5. As an example for their children	19.7%	35.7%	+81%


Figure 26
Most Important Reason for Wanting to Quit Cigarette Smoking
 Maryland Adults, 2008



Source: Maryland Adult Tobacco Surveys, 2008.

Why Smokers Were Unable to Quit Smoking



Smokers who had tried to quit but had not succeeded offered a variety of reasons as to why they were not successful during their most recent quit attempt. The most common reason cited was being nervous, tense, angry, frustrated, or stressed and that smoking helped address those issues. Figure 27 presents data on the top 10 reasons given in 2008 for being unsuccessful when trying to quit smoking.



Where Information Was Obtained on Quitting

Survey respondents who had attempted to quit smoking during the previous 12 months were asked to identify all sources where they received their information about smoking cessation from a comprehensive list. Over 80% of those who had tried to quit indicated that they had received or obtained information about smoking cessation. Figure 28 details both organizational and media sources for cessation information.

Figure 28
Where Smokers Who Tried to Quit During the Past 12 Months
Received Information About Quitting Cigarette Smoking
 Maryland Adults, 2008



Source: Maryland Adult Tobacco Surveys, 2008.

Selected Attitudes and Opinions About Smoking

Results of the Maryland Adult Tobacco Survey 2008 study showed that in general current smokers and non-smokers have similar attitudes and opinions regarding smoking and its effects.

Q. How addictive do you think cigarette smoking is?

<u>2008</u>	<u>Not at all Addictive</u>	<u>Moderately Addictive</u>	<u>Very Addictive</u>
Current Smoker	2.8%	15.1%	80.1%
Non-Smoker	2.5%	10.3%	84.1%

Q. Smoking light cigarettes is safer than smoking regular cigarettes?

<u>2008</u>	<u>Strongly Agree</u>	<u>Moderately Agree</u>	<u>Moderately Disagree</u>	<u>Strongly Disagree</u>
Current Smokers	5.6%	11.2%	14.9%	65.5%
Non-Smokers	4.0%	9.3%	13.4%	67.2%

Q. If a person smoked a pack of cigarettes every day for more than 20 years, there is LITTLE health benefit to quitting smoking?

<u>2008</u>	<u>Strongly Agree</u>	<u>Somewhat Agree</u>	<u>Somewhat Disagree</u>	<u>Strongly Disagree</u>
Current Smokers	18.3%	13.6%	17.1%	48.9%
Non-Smokers	13.7%	8.9%	13.6%	61.3%

Q. Do you agree or disagree with the following statement: "Smoke from other people's cigarettes is harmful to children." Would you say you...?

<u>2008</u>	<u>Strongly Disagree</u>	<u>Somewhat Disagree</u>	<u>Somewhat Agree</u>	<u>Strongly Agree</u>
Current Smokers	3.5%	4.5%	16.0%	75.0%
Non-Smokers	3.0%	1.2%	8.1%	86.9%

Q. How important is it to prevent sales of tobacco products to youth under 18?


<u>2008</u>	<u>Not Important</u>	<u>Somewhat Important</u>	<u>Very Important</u>
Current Smokers	4.3%	14.7%	80.2%
Non-Smokers	2.1%	8.4%	88.8%

Source for all questions and responses: Maryland Adult Tobacco Survey, 2008.

Part 4 — Secondhand Smoke


As with the previous section, the most recent data available on Maryland adult exposure to secondhand smoke is from the 2008 MATS. Thus, the content that follows is derived from those data.

The critical program elements identified in the CDC Best Practice Recommendations as essential to changing the behaviors of adults with regard to exposure to secondhand smoke include: dissemination of information about the dangers that smoke from tobacco products poses to those who inhale it secondhand, creation and enforcement of smoke-free policies, and changes in attitudes and beliefs about secondhand smoke. Together, these elements can create a social norm where involuntary exposure to secondhand smoke is not an acceptable option.





In 1993, Maryland adopted regulations that banned smoking in places of indoor employment and public spaces, with the exception of bars and restaurants with licenses to sell alcoholic beverages. Under this regulatory scheme, an estimated 85% of Maryland employees were protected from secondhand smoke in the workplace. Effective February 1, 2008, Maryland's comprehensive Clean Indoor Air Act banned smoking in indoor workplaces (including bars and all restaurants) and almost all other indoor public spaces. This legislation protects almost 100% of Maryland workers from being exposed to secondhand smoke in indoor work settings.

In the fall of 2008, 94.6% of adult respondents who were employed primarily in indoor work environments reported compliance in the workplace within the previous 30 days. There is no significant variation by race or ethnicity in this measure: 94.1% among Black/African-Americans; 93.6% among Hispanics; and 94.6% among Whites.



Underage Youth Exposure

Underage youth exposure to secondhand smoke can occur at home, in other indoor locations, or in the car. In addition to the Clean Indoor Air Act’s mandatory smoking ban in almost all indoor public areas, local health department tobacco control programs have encouraged voluntary in-home and in-car prohibitions on smoking to reduce this exposure. There was a significant increase in the percentage of underage youth living in non-smoking households who reported that smoking was not allowed in their home. Between 2008 and 2010, there were significant increases in the percentage of underage youth not exposed to secondhand smoke in a room or car.



1. US Centers for Disease Control and Prevention. Smoking-caused deaths in Maryland: smoking-attributable mortality, morbidity, and economic costs (SAMMEC) online application [Internet]. Atlanta (GA): US Centers for Disease Control and Prevention; 2004 [cited 2010 Jul 8]. Available from: https://apps/nccd.cdc.gov/sammec/five_yr_sam.asp?year=2004&state=MD
2. Hyland A, Li Q, Bauer J, Giovino GA, Yang Y, Cummings KM. Cigarette smoking attributable morbidity by state. Report by Roswell Park Cancer Institute. Buffalo (NY): Roswell Park Cancer Institute; 2003 Sep 5 [cited 2010 Jul 13]. Available from: http://roswell.tobaccodocuments.org/morbidity/morbidity_by_state.htm
3. Campaign for Tobacco-Free Kids. State-Specific Tobacco Company Marketing Expenditures 1998-2008. Washington DC, 2011 [cited 2011 Sept 6]. Available from: <http://www.tobaccofreekids.org/research/factsheets/pdf/0271.pdf>

Appendix A

Youth Prevalence, Initiation, and Cessation

	<u>Page</u>
Maryland	36
Allegany	38
Anne Arundel	40
Baltimore City	42
Baltimore County	44
Calvert	46
Caroline	48
Carroll	50
Cecil	52
Charles	54
Dorchester	56
Frederick	58
Garrett	60
Harford	62
Howard	64
Kent	66
Montgomery	68
Prince George's	70
Queen Anne's	72
Somerset	74
St. Mary's	76
Talbot	78
Washington	80
Wicomico	82
Worcester	84



Maryland

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program
Glossary of Terms

The glossary of terms is provided to assist readers in interpreting the use of the following analytical concepts:

Current tobacco use –

Students who reported using any tobacco product (cigarette, cigar, smokeless tobacco, pipe, bidis, or kreteks) in the past 30 days.

Current use of bidis, kreteks, or pipes –

In 2010, due to limitations on the number of questions that could be included in the survey, individual questions about the use of each of these tobacco products were not asked. Instead, a single question asked about the use of these products collectively, i.e., use of any one or a combination of these products during the previous 30 days. Using this question form supports the reported estimates of “any tobacco use” during the previous 30 days, but not the use of any of these individual tobacco products during the past 30 days.

Initiation in the past year –

Students who reported that they began using any tobacco product in the past 12 months (cigarette, cigar, smokeless tobacco, pipe, bidis, or kreteks).

Note: This concept captures general initiation of any new tobacco products. It does not necessarily capture first initiation of tobacco use in the past 12 months.

Cessation in the past year –

Students who quit smoking cigarettes in the past 12 months.

Note: Due to limitations of the survey data, cessation of tobacco products cannot be calculated. Thus, this concept includes only cessation of cigarettes in the past 12 months.



Maryland

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	21.4% ± 1.2% 87,963	18.5% ± 1.0%	15.6% ± 0.6%	16.2% ± 0.5%	*17.1% ± 0.5% 74,173
Any Tobacco Product (Minority Youth)	18.9% ± 1.4% 33,913	16.9% ± 1.2%	14.4% ± 0.8%	16.3% ± 0.7%	17.9% ± 0.7% 43,110
Cigarette	15.7% ± 1.1% 61,808	12.5% ± 0.8%	10.0% ± 0.5%	10.2% ± 0.4%	*9.6% ± 0.4% 40,945
Cigar	8.8% ± 0.6% 35,268	7.5% ± 0.5%	6.5% ± 0.4%	10.6% ± 0.4%	9.5% ± 0.4% 40,752
Pipe (tobacco)	3.9% ± 0.4% 15,670	4.9% ± 0.4%	4.7% ± 0.3%	5.9% ± 0.4%	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.3% ± 0.5% 21,077	5.4% ± 0.4%	4.8% ± 0.4%	4.0% ± 0.3%	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.8% ± 0.3% 15,285	4.3% ± 0.4%	5.1% ± 0.4%	3.6% ± 0.3%	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.5% ± 0.3% 14,109	3.7% ± 0.3%	3.6% ± 0.3%	3.7% ± 0.2%	3.3% ± 0.2% 13,746

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	19.5% ± 1.0% 80,050	16.2% ± 0.8%	15.0% ± 0.6%	14.8% ± 0.4%	18.1% ± 0.5% 78,621
Minority Youth	16.7% ± 1.0% 29,886	13.7% ± 0.9%	12.4% ± 0.7%	13.2% ± 0.5%	*18.9% ± 0.6% 45,605

CESSATION OF CIGARETTE USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	35.1% ± 1.9% 16,304	36.0% ± 1.7%	40.2% ± 1.9%	38.0% ± 1.8%	*45.0% ± 1.8% 12,442
Minority Youth	44.5% ± 4.0% 6,629	43.4% ± 3.1%	47.7% ± 3.1%	44.9% ± 2.7%	49.7% ± 2.5% 7,402

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	7.2% ± 0.9% 13,134	5.2% ± 0.5% 10,110	3.7% ± 0.5% 7,029	3.5% ± 0.4% 6,388	*3.5% ± 0.4% 6,294
MS Females	7.2% ± 1.1% 6,319	5.1% ± 0.7% 4,798	3.2% ± 0.6% 2,936	2.9% ± 0.5% 2,584	*3.1% ± 0.5% 2,929
MS Males	7.2% ± 1.2% 6,679	5.4% ± 0.7% 5,233	4.2% ± 0.7% 4,017	4.1% ± 0.6% 3,793	*3.8% ± 0.5% 3,328
MS Minorities	6.5% ± 1.3% 5,221	5.3% ± 0.8% 5,057	3.9% ± 0.7% 3,924	3.9% ± 0.5% 3,625	*4.1% ± 0.6% 4,179
High School (HS)	23.0% ± 0.9% 48,674	18.7% ± 0.8% 41,822	14.7% ± 0.7% 36,720	15.3% ± 0.5% 36,190	*14.1% ± 0.6% 34,651
HS Females	23.4% ± 1.3% 24,702	18.7% ± 1.0% 21,121	13.7% ± 0.9% 17,315	13.2% ± 0.6% 15,589	*11.7% ± 0.7% 14,966
HS Males	22.4% ± 1.2% 23,551	18.4% ± 1.0% 20,113	15.6% ± 0.8% 19,299	17.4% ± 0.7% 20,444	*16.6% ± 0.8% 19,580
HS Minorities	16.0% ± 1.3% 14,204	13.5% ± 0.9% 13,884	11.1% ± 0.8% 14,390	12.7% ± 0.6% 14,782	*12.2% ± 0.6% 16,446

Maryland - Current Cigarette Smoking

(Underage Middle & High School Youth, By Grade)





Allegany County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	30.1% ± 3.7% 1,688	26.8% ± 3.8% 1,468	22.5% ± 3.0% 1,145	20.1% ± 2.6% 964	*19.1% ± 2.5% 874
Any Tobacco Product (Minority Youth)	30.8% ± 7.3% 154	33.7% ± 7.7% 186	29.1% ± 6.3% 174	32.1% ± 5.8% 160	25.1% ± 5.3% 157
Cigarette	24.6% ± 3.4% 1,320	21.3% ± 3.6% 1,132	16.6% ± 2.5% 826	14.9% ± 2.2% 705	*13.5% ± 2.1% 606
Cigar	9.3% ± 1.5% 511	8.0% ± 1.8% 433	9.4% ± 1.7% 470	10.9% ± 1.9% 520	9.1% ± 1.7% 410
Pipe (tobacco)	4.7% ± 1.4% 260	4.4% ± 1.2% 238	4.1% ± 1.0% 205	5.3% ± 1.3% 252	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	4.4% ± 1.1% 21,077	5.6% ± 1.3% 22,554	4.1% ± 1.1% 21,247	3.5% ± 0.9% 166	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.3% ± 1.0% 181	3.6% ± 1.1% 197	4.1% ± 1.1% 208	3.1% ± 0.8% 145	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	9.1% ± 1.6% 503	8.6% ± 2.0% 460	6.1% ± 1.3% 305	7.0% ± 1.3% 327	6.6% ± 1.3% 292

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	27.4% ± 2.8% 1,536	21.9% ± 2.6% 1,204	22.6% ± 2.8% 1,151	19.5% ± 2.3% 933	*19.1% ± 2.3% 876
Minority Youth	24.5% ± 5.6% 122	22.7% ± 6.4% 125	17.8% ± 4.5% 107	23.6% ± 5.3% 117	24.8% ± 4.6% 155

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	31.2% ± 4.7% 336	26.2% ± 6.4% 205	30.1% ± 5.5% 165	30.6% ± 6.6% 174	34.8% ± 7.8% 129
Minority Youth	20.5% ± 15.2% 23	Not Available	Not Available	25.1% ± 11.6% 25	18.6% ± 11.2% 12

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.9% ± 2.9% 229	8.3% ± 2.3% 195	6.9% ± 2.5% 140	3.9% ± 1.4% 77	6.4% ± 2.2% 123
MS Females	8.5% ± 3.2% 94	9.4% ± 3.2% 105	7.7% ± 4.4% 76	3.8% ± 1.9% 37	6.0% ± 2.3% 60
MS Males	11.3% ± 3.9% 135	7.3% ± 2.8% 89	6.2% ± 2.9% 64	4.1% ± 1.5% 41	6.8% ± 3.0% 62
MS Minorities	15.3% ± 7.3% 40	11.9% ± 6.1% 35	7.0% ± 4.7% 20	10.7% ± 5.1% 25	12.6% ± 6.7% 40
High School (HS)	35.6% ± 3.7% 1,092	31.6% ± 4.9% 937	23.4% ± 2.5% 686	22.8% ± 3.6% 627	*18.7% ± 3.0% 483
HS Females	37.7% ± 4.5% 575	34.8% ± 5.9% 525	24.8% ± 3.5% 347	22.5% ± 4.4% 299	*16.5% ± 3.2% 223
HS Males	33.2% ± 4.4% 505	28.0% ± 5.7% 405	22.0% ± 3.1% 337	23.0% ± 4.4% 328	*21.1% ± 4.7% 258
HS Minorities	37.0% ± 10.1% 78	44.9% ± 13.9% 95	26.5% ± 6.3% 72	40.5% ± 7.8% 101	28.6% ± 6.7% 82

Allegany County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Anne Arundel County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	24.8% ± 4.5% 9,349	20.4% ± 3.4% 8,030	16.5% ± 2.1% 6,465	18.4% ± 2.1% 6,910	18.2% ± 3.2% 7,016
Any Tobacco Product (Minority Youth)	24.9% ± 5.5% 2,254	18.7% ± 4.7% 2,044	16.9% ± 3.0% 2,256	21.4% ± 3.3% 2,559	21.9% ± 3.0% 3,048
Cigarette	19.8% ± 4.0% 7,203	14.7% ± 3.0% 5,588	11.1% ± 1.6% 4,253	12.2% ± 1.5% 4,517	*10.4% ± 2.2% 3,928
Cigar	10.7% ± 2.3% 3,946	9.0% ± 1.9% 3,445	7.3% ± 1.1% 2,773	<u>12.0%</u> ± 1.5% 4,431	9.6% ± 2.2% 3,664
Pipe (tobacco)	4.4% ± 1.1% 1,620	5.5% ± 1.0% 2,100	5.4% ± 1.1% 2,096	6.6% ± 1.1% 2,424	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.2% ± 1.2% 1,895	5.2% ± 1.0% 1,989	5.3% ± 1.1% 2,052	4.4% ± 0.8% 1,615	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.7% ± 1.1% 1,730	4.7% ± 1.0% 1,803	5.0% ± 1.0% 1,938	3.9% ± 0.8% 1,441	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.5% ± 1.1% 1,306	4.0% ± 0.9% 1,499	3.9% ± 0.8% 1,500	3.5% ± 0.7% 1,256	3.3% ± 0.8% 1,222

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	23.5% ± 3.2% 8,883	18.1% ± 2.7% 7,116	16.5% ± 1.9% 6,485	15.4% ± 1.5% 5,807	18.0% ± 3.2% 6,931
Minority Youth	21.8% ± 3.9% 1,963	15.3% ± 3.2% 1,671	14.9% ± 2.3% 1,989	15.2% ± 2.2% 1,812	20.0% ± 3.4% 2,792

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	31.0% ± 4.7% 1,684	32.3% ± 5.6% 1,159	40.3% ± 5.1% 1,144	34.6% ± 4.9% 1,234	41.3% ± 6.3% 1,053
Minority Youth	37.7% ± 9.5% 410	28.7% ± 10.8% 165	44.8% ± 9.7% 408	37.9% ± 9.3% 483	44.3% ± 9.5% 402

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk “*”.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.4% ± 3.1% 1,556	6.0% ± 2.0% 1,055	3.7% ± 1.2% 602	4.4% ± 2.0% 703	*3.3% ± 1.5% 540
MS Females	11.1% ± 4.3% 873	6.6% ± 2.6% 568	3.3% ± 1.7% 263	4.2% ± 2.9% 333	*3.4% ± 2.0% 291
MS Males	7.7% ± 3.0% 667	5.3% ± 2.4% 465	4.0% ± 1.7% 330	4.6% ± 1.8% 370	3.1% ± 1.5% 249
MS Minorities	9.4% ± 4.6% 343	5.6% ± 3.0% 266	4.5% ± 2.4% 243	7.4% ± 3.6% 381	5.0% ± 2.1% 308
High School (HS)	28.5% ± 3.2% 5,647	22.5% ± 2.3% 4,533	16.7% ± 1.7% 3,651	18.3% ± 2.1% 3,814	*15.9% ± 1.9% 3,388
HS Females	28.4% ± 3.7% 2,769	23.2% ± 2.8% 2,378	16.2% ± 2.2% 1,797	16.0% ± 2.3% 1,674	*13.7% ± 2.1% 1,496
HS Males	28.6% ± 4.2% 2,864	21.1% ± 3.0% 2,131	17.0% ± 2.1% 1,826	20.3% ± 2.6% 2,116	*18.2% ± 2.5% 1,892
HS Minorities	23.8% ± 5.1% 1,152	15.9% ± 2.9% 875	15.6% ± 2.4% 1,149	16.0% ± 2.6% 1,022	*15.3% ± 2.2% 1,132

Anne Arundel County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Baltimore City

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	20.3% ± 3.0% 8,961	18.5% ± 2.6% 8,321	15.5% ± 2.4% 6,350	<u>20.7%</u> ± 2.1% 6,507	21.1% ± 2.9% 8,192
Any Tobacco Product (Minority Youth)	19.5% ± 3.1% 7,450	17.0% ± 2.5% 6,757	15.0% ± 2.4% 5,744	<u>20.4%</u> ± 4.6% 5,810	20.8% ± 2.9% 7,472
Cigarette	10.8% ± 2.1% 4,423	9.5% ± 1.6% 3,954	7.7% ± 1.4% 2,985	10.2% ± 1.3% 3,072	9.5% ± 1.7% 3,641
Cigar	8.8% ± 1.8% 3,638	7.7% ± 1.6% 3,164	6.3% ± 1.1% 2,357	<u>16.2%</u> ± 2.1% 4,796	*13.5% ± 2.1% 5,067
Pipe (tobacco)	4.2% ± 1.2% 1,724	6.6% ± 1.4% 2,707	5.4% ± 1.6% 2,118	<u>9.8%</u> ± 1.8% 2,767	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	6.5% ± 2.0% 2,600	8.1% ± 1.9% 3,350	6.1% ± 1.5% 2,372	7.2% ± 1.3% 2,017	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.5% ± 1.4% 1,830	6.3% ± 1.7% 2,611	6.5% ± 1.6% 2,494	6.1% ± 1.3% 1,711	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.6% ± 1.1% 1,490	4.3% ± 1.2% 1,768	3.1% ± 0.9% 1,187	4.2% ± 0.9% 1,141	2.9% ± 0.9% 1,011

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	19.2% ± 2.9% 8,425	14.5% ± 1.7% 6,532	14.2% ± 1.8% 5,786	16.9% ± 1.4% 5,294	<u>21.7%</u> ± 2.3% 8,423
Minority Youth	17.4% ± 2.1% 6,597	13.5% ± 1.8% 5,369	13.9% ± 1.8% 5,312	16.5% ± 1.4% 4,686	<u>21.6%</u> ± 2.3% 7,758

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	45.8% ± 7.3% 1,978	44.8% ± 8.7% 1,185	47.8% ± 9.4% 1,027	43.7% ± 5.2% 1,282	43.5% ± 6.2% 1,088
Minority Youth	48.8% ± 9.8% 1,538	48.5% ± 10.9% 985	51.1% ± 10.5% 940	46.7% ± 5.5% 1,219	45.1% ± 6.7% 987

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.0% ± 3.0% 1,819	7.7% ± 1.9% 1,524	6.6% ± 2.5% 1,153	7.0% ± 1.7% 972	5.8% ± 1.0% 950
MS Females	8.9% ± 3.9% 874	6.4% ± 1.7% 628	5.5% ± 3.1% 468	5.5% ± 2.2% 383	5.5% ± 1.2% 454
MS Males	9.2% ± 4.7% 945	9.1% ± 3.6% 896	7.3% ± 3.5% 657	8.5% ± 2.1% 585	6.0% ± 1.7% 476
MS Minorities	7.7% ± 2.9% 1,304	7.2% ± 2.2% 1,273	5.5% ± 2.3% 894	6.7% ± 1.5% 839	5.3% ± 0.9% 794
High School (HS)	12.5% ± 2.9% 2,605	11.1% ± 2.0% 2,430	8.6% ± 1.3% 1,832	12.9% ± 2.0% 2,101	12.3% ± 2.4% 2,692
HS Females	12.0% ± 4.1% 1,278	9.7% ± 2.6% 1,152	6.9% ± 1.5% 791	10.0% ± 1.9% 840	7.6% ± 1.9% 861
HS Males	13.1% ± 4.5% 1,316	12.5% ± 3.0% 1,235	10.4% ± 2.2% 1,032	15.7% ± 2.7% 1,243	17.1% ± 2.9% 1,809
HS Minorities	11.2% ± 2.8% 2,063	8.6% ± 2.0% 1,647	8.1% ± 1.4% 1,634	11.7% ± 2.1% 1,730	11.6% ± 2.2% 2,364

Baltimore City - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Baltimore County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	21.6% ± 3.9% 11,445	18.4% ± 4.0% 10,268	16.1% ± 1.9% 9,059	18.8% ± 2.0% 9,391	18.2% ± 3.4% 9,519
Any Tobacco Product (Minority Youth)	18.6% ± 3.5% 3,388	15.2% ± 3.7% 3,575	13.2% ± 1.9% 3,819	<u>17.2%</u> ± 1.8% 4,274	17.7% ± 3.5% 4,919
Cigarette	15.8% ± 3.6% 8,046	12.8% ± 3.2% 6,815	10.4% ± 1.5% 5,628	12.1% ± 1.7% 5,866	10.8% ± 2.5% 5,548
Cigar	9.0% ± 2.4% 4,650	8.1% ± 1.9% 4,345	6.6% ± 1.0% 3,565	<u>12.7%</u> ± 1.5% 6,172	11.1% ± 2.5% 5,730
Pipe (tobacco)	3.4% ± 1.1% 1,799	4.6% ± 1.2% 2,435	4.6% ± 0.8% 2,547	<u>7.6%</u> ± 1.5% 3,634	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.3% ± 1.2% 2,731	4.9% ± 1.3% 2,631	4.8% ± 0.9% 2,633	4.5% ± 0.9% 2,150	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.0% ± 0.9% 2,103	3.9% ± 1.2% 2,086	5.2% ± 0.9% 2,866	4.7% ± 1.1% 2,233	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	2.6% ± 0.7% 1,358	2.9% ± 0.8% 1,565	3.4% ± 0.6% 1,860	3.9% ± 1.0% 1,808	3.0% ± 1.0% 1,477

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	18.5% ± 3.4% 9,829	16.9% ± 2.7% 9,431	15.4% ± 1.6% 8,663	15.8% ± 1.9% 7,861	19.5% ± 3.2% 10,181
Minority Youth	15.2% ± 2.9% 2,775	12.2% ± 2.8% 2,890	11.1% ± 1.6% 3,207	12.3% ± 1.9% 3,038	<u>19.9%</u> ± 3.4% 5,541

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	34.2% ± 6.7% 2,014	32.5% ± 5.6% 1,501	33.5% ± 4.6% 1,243	32.6% ± 6.1% 1,534	42.4% ± 7.3% 1,550
Minority Youth	54.1% ± 9.6% 667	45.4% ± 10.9% 623	47.1% ± 8.9% 584	42.0% ± 7.7% 766	50.1% ± 10.7% 809

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	6.5% ± 3.3% 1,536	4.6% ± 1.9% 1,121	3.0% ± 1.2% 696	4.6% ± 2.0% 996	3.0% ± 1.0% 670
MS Females	8.2% ± 3.9% 955	4.9% ± 2.3% 586	1.8% ± 1.1% 198	4.1% ± 2.5% 444	*2.3% ± 1.2% 258
MS Males	4.9% ± 3.3% 574	4.4% ± 1.9% 536	4.2% ± 1.8% 497	5.0% ± 2.0% 552	3.8% ± 2.2% 412
MS Minorities	3.8% ± 1.8% 321	4.7% ± 2.2% 528	2.6% ± 1.5% 316	3.4% ± 1.2% 386	3.5% ± 1.3% 423
High School (HS)	23.7% ± 3.0% 6,510	19.7% ± 3.4% 5,694	15.8% ± 1.6% 4,933	18.3% ± 2.6% 4,870	*16.7% ± 2.5% 4,878
HS Females	25.6% ± 4.5% 3,501	20.3% ± 3.7% 2,973	15.1% ± 2.0% 2,399	16.1% ± 3.0% 2,151	*15.1% ± 2.8% 2,262
HS Males	21.5% ± 3.1% 2,932	18.9% ± 4.2% 2,655	16.5% ± 2.1% 2,526	20.4% ± 2.6% 2,711	18.4% ± 3.0% 2,609
HS Minorities	16.2% ± 2.7% 1,445	11.9% ± 4.1% 1,315	9.8% ± 1.6% 1,496	12.8% ± 1.6% 1,629	*11.5% ± 1.8% 1,733

Baltimore County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Calvert County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	24.2% ± 2.7% 1,932	22.5% ± 3.8% 1,979	17.2% ± 2.4% 1,677	14.7% ± 1.7% 1,382	*17.4% ± 2.2% 1,615
Any Tobacco Product (Minority Youth)	24.8% ± 4.4% 433	27.4% ± 6.8% 527	21.0% ± 3.4% 519	17.7% ± 3.4% 350	18.3% ± 3.0% 415
Cigarette	19.6% ± 2.7% 1,502	17.9% ± 3.5% 1,505	13.3% ± 2.0% 1,270	10.6% ± 1.4% 985	*11.6% ± 1.7% 1,060
Cigar	10.0% ± 1.8% 786	10.1% ± 2.3% 856	6.4% ± 1.2% 612	7.2% ± 1.1% 671	7.8% ± 1.3% 716
Pipe (tobacco)	3.6% ± 0.9% 286	6.5% ± 1.9% 551	3.8% ± 0.9% 367	3.7% ± 0.8% 344	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	3.9% ± 0.9% 306	5.5% ± 1.5% 467	3.8% ± 1.0% 363	2.5% ± 0.7% 232	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.5% ± 0.8% 274	5.4% ± 2.1% 455	4.1% ± 1.0% 391	2.3% ± 0.6% 211	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	2.8% ± 0.7% 220	4.0% ± 1.4% 340	3.4% ± 0.8% 320	4.1% ± 0.8% 374	3.7% ± 0.9% 338

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	23.3% ± 2.1% 1,857	19.5% ± 2.9% 1,717	17.5% ± 2.1% 1,707	14.4% ± 1.5% 1,353	*18.4% ± 2.1% 1,708
Minority Youth	20.5% ± 3.4% 358	19.4% ± 4.4% 373	16.2% ± 2.6% 398	12.9% ± 2.6% 256	20.9% ± 3.2% 472

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	31.2% ± 4.8% 367	26.7% ± 5.6% 275	36.1% ± 5.6% 311	38.5% ± 5.9% 299	39.2% ± 5.8% 279
Minority Youth	27.6% ± 9.4% 61	29.7% ± 13.3% 62	37.9% ± 11.8% 80	36.7% ± 10.7% 72	39.0% ± 11.2% 67

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '**’.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.6% ± 3.4% 337	6.1% ± 2.4% 238	3.8% ± 1.5% 152	2.9% ± 1.2% 114	*3.0% ± 1.2% 115
MS Females	12.2% ± 5.0% 199	5.3% ± 2.9% 99	3.8% ± 1.9% 75	2.7% ± 1.5% 52	*3.3% ± 1.9% 64
MS Males	7.2% ± 3.3% 137	6.8% ± 2.8% 139	3.7% ± 2.1% 77	3.1% ± 1.8% 62	2.7% ± 1.5% 51
MS Minorities	11.8% ± 6.6% 97	7.4% ± 3.9% 69	7.3% ± 3.4% 74	5.3% ± 3.6% 45	4.2% ± 2.2% 40
High School (HS)	28.3% ± 3.6% 1,164	28.1% ± 5.8% 1,267	20.3% ± 2.1% 1,119	16.2% ± 2.2% 871	*17.7% ± 2.3% 945
HS Females	29.2% ± 4.3% 604	26.1% ± 6.5% 610	20.8% ± 2.6% 587	15.7% ± 2.7% 421	*15.7% ± 2.7% 435
HS Males	27.3% ± 4.3% 557	30.4% ± 7.2% 657	19.8% ± 3.0% 532	16.7% ± 2.8% 445	19.9% ± 3.2% 509
HS Minorities	23.2% ± 5.2% 196	33.5% ± 10.5% 290	19.8% ± 3.8% 268	17.4% ± 3.1% 193	19.6% ± 3.6% 247

Calvert County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Caroline County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	31.3% ± 3.9% 870	25.7% ± 2.4% 747	22.7% ± 2.7% 663	21.6% ± 2.5% 599	*20.9% ± 2.7% 548
Any Tobacco Product (Minority Youth)	29.6% ± 5.7% 205	24.5% ± 4.6% 184	24.3% ± 4.0% 196	21.5% ± 3.5% 160	24.1% ± 4.1% 206
Cigarette	25.8% ± 3.6% 691	<u>19.8%</u> ± 2.0% 554	16.9% ± 2.4% 842	17.0% ± 2.3% 460	*14.3% ± 2.3% 366
Cigar	13.2% ± 2.3% 362	10.0% ± 1.4% 285	8.6% ± 1.5% 246	<u>12.0%</u> ± 1.8% 328	10.5% ± 1.9% 273
Pipe (tobacco)	5.0% ± 1.5% 138	5.0% ± 1.2% 143	4.8% ± 1.1% 138	6.2% ± 1.2% 167	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.5% ± 1.3% 149	4.8% ± 1.3% 138	3.8% ± 1.0% 110	3.8% ± 1.0% 104	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	5.9% ± 1.5% 162	4.1% ± 1.0% 117	4.1% ± 1.1% 119	3.4% ± 1.0% 93	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	6.3% ± 1.8% 172	5.0% ± 1.2% 140	5.6% ± 1.1% 161	6.0% ± 1.3% 159	6.6% ± 1.4% 167

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	25.2% ± 3.0% 700	22.7% ± 2.2% 661	23.5% ± 2.5% 689	19.9% ± 2.1% 551	19.8% ± 2.6% 517
Minority Youth	20.1% ± 5.1% 139	22.8% ± 3.4% 172	20.2% ± 3.3% 163	15.7% ± 3.0% 116	<u>23.2%</u> ± 4.1% 198

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	26.0% ± 6.4% 124	28.2% ± 5.1% 96	35.1% ± 6.5% 117	31.7% ± 5.5% 109	*41.4% ± 8.3% 96
Minority Youth	22.5% ± 11.2% 24	41.4% ± 13.8% 26	36.2% ± 12.0% 29	45.9% ± 12.3% 38	36.9% ± 13.6% 23

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '**’.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	13.5% ± 3.9% 165	10.6% ± 2.5% 141	7.5% ± 1.5% 89	6.0% ± 2.1% 70	*3.4% ± 1.4% 38
MS Females	11.7% ± 6.3% 69	10.5% ± 3.2% 68	7.9% ± 2.5% 44	4.7% ± 2.4% 26	*2.9% ± 1.4% 16
MS Males	15.4% ± 4.4% 96	10.7% ± 2.9% 73	7.2% ± 2.0% 45	7.3% ± 2.7% 44	*4.0% ± 2.1% 22
MS Minorities	13.9% ± 6.0% 46	11.8% ± 4.4% 43	9.8% ± 3.3% 31	6.1% ± 3.0% 20	*3.2% ± 1.9% 13
High School (HS)	36.0% ± 5.0% 527	28.3% ± 3.5% 413	23.7% ± 2.9% 393	25.1% ± 3.5% 390	*22.6% ± 3.4% 328
HS Females	35.5% ± 5.9% 249	28.9% ± 4.2% 207	23.1% ± 3.7% 191	22.5% ± 4.2% 172	*19.1% ± 3.5% 149
HS Males	36.3% ± 6.0% 274	27.1% ± 5.0% 198	24.0% ± 3.6% 199	27.6% ± 4.8% 217	26.7% ± 5.1% 178
HS Minorities	30.0% ± 7.8% 96	22.5% ± 6.7% 76	21.4% ± 4.5% 93	19.9% ± 4.5% 76	24.6% ± 5.5% 105

Caroline County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Carroll County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	21.1% ± 3.4% 3,014	18.9% ± 3.4% 2,842	17.0% ± 2.2% 2,764	15.3% ± 1.6% 2,372	*15.0% ± 1.6% 2,263
Any Tobacco Product (Minority Youth)	34.8% ± 7.7% 463	26.9% ± 6.1% 466	23.0% ± 4.2% 436	23.9% ± 4.1% 327	*18.0% ± 3.0% 319
Cigarette	16.4% ± 3.0% 2,263	13.9% ± 2.9% 2,024	12.1% ± 1.7% 1,932	11.2% ± 1.2% 1,709	*9.5% ± 1.2% 1,419
Cigar	7.6% ± 1.9% 1,076	7.9% ± 1.7% 1,169	7.9% ± 1.2% 1,263	7.8% ± 1.1% 1,196	7.2% ± 1.0% 1,084
Pipe (tobacco)	3.1% ± 0.8% 438	4.7% ± 1.3% 688	3.4% ± 0.6% 551	3.4% ± 0.7% 525	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	4.8% ± 1.1% 674	4.9% ± 1.4% 728	2.9% ± 0.6% 477	2.5% ± 0.6% 384	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.3% ± 0.8% 469	4.0% ± 1.2% 595	3.3% ± 0.6% 527	1.9% ± 0.4% 292	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.2% ± 0.9% 448	4.0% ± 1.1% 587	3.8% ± 0.7% 614	4.7% ± 0.8% 707	4.3% ± 0.7% 638

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	17.7% ± 2.3% 2,522	16.2% ± 2.6% 2,433	18.0% ± 2.3% 2,940	15.7% ± 1.6% 2,425	16.2% ± 1.7% 2,451
Minority Youth	22.3% ± 5.3% 293	15.4% ± 5.4% 267	15.7% ± 3.5% 298	14.2% ± 3.6% 194	17.9% ± 3.1% 318

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	34.2% ± 6.1% 558	27.8% ± 8.2% 352	36.0% ± 5.1% 385	30.6% ± 5.0% 410	43.0% ± 6.3% 372
Minority Youth	26.0% ± 15.2% 49	Not Available	41.3% ± 17.0% 298	22.3% ± 11.4% 33	45.9% ± 15.4% 46

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '**’.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	4.9% ± 2.8% 302	3.1% ± 1.6% 208	3.5% ± 1.3% 233	1.6% ± 0.8% 103	1.8% ± 1.0% 109
MS Females	4.5% ± 3.2% 134	3.3% ± 2.4% 108	3.4% ± 1.5% 111	1.8% ± 1.8% 54	1.6% ± 1.1% 50
MS Males	5.2% ± 2.9% 168	2.5% ± 1.6% 86	3.6% ± 2.0% 122	1.5% ± 0.8% 49	2.0% ± 1.3% 59
MS Minorities	11.2% ± 9.9% 71	8.0% ± 5.5% 65	4.2% ± 4.1% 36	6.0% ± 3.9% 38	3.2% ± 2.3% 27
High School (HS)	25.9% ± 3.7% 1,960	23.1% ± 4.2% 1,816	18.1% ± 1.8% 1,699	17.8% ± 1.9% 1,606	*14.8% ± 1.7% 1,310
HS Females	26.2% ± 4.8% 963	23.4% ± 5.2% 916	17.7% ± 2.3% 808	15.5% ± 2.2% 693	*13.2% ± 2.0% 593
HS Males	25.6% ± 3.9% 975	22.3% ± 4.6% 860	18.5% ± 2.2% 887	20.0% ± 2.6% 902	*16.5% ± 2.4% 717
HS Minorities	42.8% ± 9.0% 275	28.6% ± 7.4% 223	20.9% ± 4.7% 191	25.4% ± 4.5% 173	*17.6% ± 3.7% 154

Carroll County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Cecil County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	26.9% ± 3.7% 2,016	22.9% ± 2.9% 1,840	21.8% ± 2.7% 1,914	20.4% ± 1.8% 1,721	*20.2% ± 2.0% 1,658
Any Tobacco Product (Minority Youth)	29.2% ± 6.5% 274	25.4% ± 6.4% 287	26.1% ± 4.1% 462	22.7% ± 3.5% 321	20.4% ± 3.2% 325
Cigarette	22.3% ± 3.4% 1,613	17.6% ± 2.7% 1,347	16.0% ± 2.4% 1,360	15.0% ± 1.5% 1,245	*13.8% ± 1.5% 1,114
Cigar	9.2% ± 1.8% 679	7.3% ± 1.5% 575	9.7% ± 1.4% 829	12.3% ± 1.5% 1,025	10.2% ± 1.4% 831
Pipe (tobacco)	3.3% ± 0.9% 248	4.2% ± 1.3% 327	5.1% ± 0.9% 442	5.9% ± 1.2% 490	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	6.2% ± 1.4% 454	6.0% ± 1.2% 474	4.0% ± 1.0% 426	3.8% ± 0.9% 309	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	2.8% ± 0.7% 208	4.2% ± 1.1% 329	5.0% ± 1.0% 434	3.4% ± 0.9% 283	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.3% ± 0.9% 242	3.8% ± 0.9% 292	5.6% ± 1.0% 484	5.7% ± 0.9% 462	5.3% ± 1.1% 421

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	23.3% ± 2.9% 1,748	18.9% ± 2.6% 1,518	20.8% ± 2.5% 1,831	19.3% ± 1.6% 1,630	21.1% ± 2.1% 1,737
Minority Youth	26.4% ± 5.2% 248	21.3% ± 5.1% 238	17.7% ± 3.0% 313	17.9% ± 3.2% 251	19.6% ± 3.1% 312

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	26.2% ± 5.0% 297	32.5% ± 5.0% 333	34.3% ± 5.5% 335	35.1% ± 4.5% 344	*42.2% ± 5.5% 317
Minority Youth	27.3% ± 13.8% 39	25.8% ± 12.6% 37	39.1% ± 12.7% 69	41.2% ± 10.2% 73	49.8% ± 10.9% 68

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*’.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	11.3% ± 2.9% 389	9.8% ± 2.7% 359	6.1% ± 2.7% 235	6.2% ± 1.9% 222	*5.0% ± 1.4% 176
MS Females	11.6% ± 4.2% 198	11.7% ± 3.6% 205	7.3% ± 4.4% 136	5.9% ± 2.2% 103	*4.9% ± 1.9% 89
MS Males	11.1% ± 3.2% 191	8.0% ± 2.9% 151	5.0% ± 2.5% 99	6.4% ± 2.5% 119	*5.1% ± 2.0% 87
MS Minorities	13.6% ± 6.4% 63	6.5% ± 3.8% 38	8.0% ± 4.5% 66	7.2% ± 3.3% 50	5.2% ± 3.1% 38
High School (HS)	32.3% ± 4.1% 1,224	24.7% ± 3.7% 988	24.0% ± 2.1% 1,125	21.8% ± 2.4% 1,023	*20.5% ± 2.1% 938
HS Females	35.3% ± 4.5% 656	28.3% ± 4.6% 581	23.4% ± 2.7% 550	21.0% ± 3.0% 495	*20.3% ± 2.8% 479
HS Males	29.4% ± 5.8% 562	21.0% ± 4.6% 403	24.5% ± 2.9% 575	22.5% ± 3.0% 522	20.7% ± 2.9% 460
HS Minorities	36.6% ± 8.9% 156	33.0% ± 9.9% 151	25.7% ± 4.6% 216	24.7% ± 4.6% 168	*19.5% ± 3.8% 164

Cecil County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Charles County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	24.6% ± 3.2% 2,957	20.4% ± 2.6% 2,572	<u>14.4%</u> ± 1.9% 2,156	16.9% ± 1.6% 2,531	*16.6% ± 1.9% 2,491
Any Tobacco Product Minority Youth)	21.9% ± 3.3% 913	17.0% ± 3.0% 955	13.8% ± 2.1% 1,264	16.6% ± 2.1% 1,491	*15.9% ± 1.8% 1,542
Cigarette	19.7% ± 3.0% 2,276	14.6% ± 2.2% 1,771	<u>9.9%</u> ± 1.7% 1,444	11.2% ± 1.3% 1,633	*9.5% ± 1.3% 1,400
Cigar	10.4% ± 2.0% 1,215	7.6% ± 1.3% 900	6.3% ± 1.0% 921	<u>11.7%</u> ± 1.4% 1,716	9.1% ± 1.2% 1,354
Pipe (tobacco)	4.8% ± 1.1% 569	5.0% ± 1.5% 589	4.0% ± 0.8% 593	5.7% ± 1.1% 831	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	6.0% ± 1.3% 694	5.2% ± 1.7% 621	4.1% ± 0.8% 608	3.5% ± 0.7% 505	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.6% ± 1.1% 544	4.6% ± 1.6% 548	4.6% ± 0.9% 680	3.4% ± 0.7% 494	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.9% ± 0.9% 242	4.1% ± 1.6% 292	3.2% ± 0.7% 484	3.4% ± 0.7% 462	3.3% ± 0.8% 471

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	21.6% ± 2.4% 2,597	18.4% ± 1.8% 2,331	<u>14.6%</u> ± 1.9% 2,187	16.0% ± 1.4% 2,398	18.9% ± 1.7% 2,834
Minority Youth	19.5% ± 3.0% 819	14.9% ± 2.4% 838	12.8% ± 2.0% 1,175	14.8% ± 1.7% 1,324	<u>18.4%</u> ± 1.8% 1,782

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	33.1% ± 5.7% 534	31.8% ± 6.6% 404	38.4% ± 5.8% 389	38.0% ± 5.3% 553	*52.3% ± 5.8% 559
Minority Youth	43.9% ± 9.7% 194	43.6% ± 14.3% 143	44.4% ± 8.4% 246	40.4% ± 7.7% 325	<u>58.7%</u> ± 7.5% 344

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.3% ± 2.8% 478	5.5% ± 1.7% 303	1.6% ± 0.8% 98	3.1% ± 1.0% 183	*3.7% ± 1.3% 224
MS Females	8.9% ± 4.2% 219	6.5% ± 2.4% 180	1.0% ± 1.2% 30	2.7% ± 1.2% 78	3.3% ± 1.6% 103
MS Males	9.7% ± 3.3% 259	4.6% ± 2.0% 123	2.2% ± 1.0% 67	3.2% ± 1.4% 99	*4.1% ± 1.9% 120
MS Minorities	8.7% ± 3.6% 166	5.1% ± 2.2% 127	1.6% ± 1.0% 63	3.2% ± 1.1% 118	*3.4% ± 1.3% 133
High School (HS)	28.0% ± 4.0% 1,797	22.0% ± 3.3% 1,469	16.0% ± 1.9% 1,347	16.8% ± 2.1% 1,450	*13.6% ± 1.8% 1,176
HS Females	26.9% ± 4.4% 860	21.2% ± 3.5% 729	15.1% ± 2.0% 656	14.2% ± 2.3% 615	*12.3% ± 2.1% 552
HS Males	28.9% ± 5.0% 923	22.7% ± 5.3% 728	16.9% ± 2.4% 691	19.4% ± 2.8% 834	*14.8% ± 2.4% 618
HS Minorities	22.5% ± 4.1% 457	14.3% ± 3.9% 401	14.0% ± 2.3% 681	14.6% ± 2.8% 735	*10.3% ± 1.7% 578

Charles County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Dorchester County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	25.5% ± 3.2% 663	22.1% ± 2.7% 587	19.7% ± 2.6% 499	18.1% ± 2.5% 427	21.6% ± 3.1% 491
Any Tobacco Product (Minority Youth)	25.3% ± 4.1% 262	22.3% ± 3.3% 236	19.6% ± 3.4% 252	21.2% ± 3.9% 236	24.6% ± 3.9% 278
Cigarette	20.0% ± 2.9% 495	15.9% ± 2.3% 405	13.7% ± 2.2% 335	13.6% ± 2.1% 312	*13.4% ± 2.2% 298
Cigar	10.7% ± 2.0% 272	9.6% ± 1.7% 247	7.8% ± 1.7% 188	10.9% ± 2.0% 248	11.2% ± 1.9% 252
Pipe (tobacco)	5.9% ± 1.6% 153	4.8% ± 1.1% 121	6.7% ± 1.4% 166	5.8% ± 1.9% 132	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	6.6% ± 1.8% 164	5.5% ± 1.3% 140	6.2% ± 1.5% 153	4.1% ± 1.2% 92	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	5.1% ± 1.4% 133	4.6% ± 1.3% 117	7.0% ± 1.7% 173	3.5% ± 1.3% 78	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.4% ± 1.3% 112	4.1% ± 1.2% 103	4.4% ± 1.1% 106	3.0% ± 1.0% 65	4.2% ± 1.2% 91

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	23.8% ± 2.5% 622	19.0% ± 2.4% 504	18.7% ± 2.4% 473	18.0% ± 2.0% 423	21.2% ± 2.9% 483
Minority Youth	23.4% ± 3.5% 243	17.3% ± 2.8% 192	16.0% ± 2.8% 206	18.8% ± 2.8% 209	23.6% ± 3.4% 268

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	36.0% ± 6.4% 126	32.5% ± 6.6% 84	41.1% ± 6.9% 93	39.8% ± 7.1% 104	38.1% ± 7.9% 69
Minority Youth	39.7% ± 13.9% 40	38.2% ± 12.4% 32	40.3% ± 11.9% 34	38.9% ± 10.3% 44	33.2% ± 9.8% 32

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk “*”.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	11.1% ± 2.9% 122	8.2% ± 2.4% 99	6.3% ± 2.1% 63	5.3% ± 1.9% 53	*5.6% ± 2.2% 53
MS Females	11.6% ± 4.0% 60	8.1% ± 2.8% 47	6.6% ± 2.4% 33	5.6% ± 2.3% 27	*4.7% ± 2.3% 24
MS Males	10.2% ± 3.7% 58	8.3% ± 3.1% 51	6.1% ± 2.7% 31	5.1% ± 2.3% 26	6.5% ± 3.2% 29
MS Minorities	12.0% ± 4.6% 54	9.4% ± 2.7% 49	6.1% ± 2.5% 32	7.4% ± 3.5% 33	5.8% ± 2.2% 28
High School (HS)	27.2% ± 3.2% 373	22.9% ± 3.7% 306	18.7% ± 2.5% 272	19.7% ± 3.5% 259	*19.4% ± 3.1% 245
HS Females	28.5% ± 4.0% 194	24.4% ± 4.4% 167	18.4% ± 3.3% 142	16.7% ± 3.7% 116	*14.6% ± 3.3% 96
HS Males	24.9% ± 4.6% 168	21.1% ± 4.8% 136	19.1% ± 4.1% 130	23.2% ± 5.5% 144	24.6% ± 4.3% 150
HS Minorities	23.3% ± 5.3% 119	19.4% ± 5.2% 102	16.0% ± 3.8% 112	19.4% ± 5.0% 122	21.9% ± 4.3% 134

Dorchester County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Frederick County Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	25.1% ± 3.2% 4,535	19.6% ± 3.0% 3,866	17.5% ± 2.3% 3,795	15.2% ± 1.5% 3,245	*15.0% ± 3.1% 3,185
Any Tobacco Product (Minority Youth)	26.3% ± 4.2% 839	26.1% ± 5.4% 955	22.0% ± 3.4% 1,216	17.1% ± 3.8% 811	*17.2% ± 3.4% 1,066
Cigarette	19.5% ± 2.8% 3,397	14.0% ± 2.5% 2,649	11.8% ± 1.8% 2,470	10.4% ± 1.2% 2,188	*9.2% ± 2.2% 1,929
Cigar	10.0% ± 1.5% 1,774	7.6% ± 1.4% 1,471	7.0% ± 1.1% 1,473	8.2% ± 1.0% 1,722	7.9% ± 2.0% 1,680
Pipe (tobacco)	4.3% ± 1.0% 779	4.9% ± 1.1% 952	5.4% ± 0.9% 1,155	4.1% ± 0.7% 851	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.2% ± 1.1% 914	5.1% ± 1.2% 989	5.3% ± 0.9% 1,141	2.8% ± 0.6% 580	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.5% ± 1.1% 808	4.2% ± 1.0% 803	5.4% ± 0.9% 1,156	2.7% ± 0.6% 547	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.7% ± 1.1% 836	5.1% ± 1.1% 980	4.9% ± 0.9% 1,030	4.6% ± 0.8% 943	4.1% ± 1.2% 850

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	22.4% ± 2.4% 4,039	17.5% ± 2.3% 3,441	16.6% ± 2.1% 3,581	14.8% ± 1.4% 3,151	16.9% ± 3.2% 3,585
Minority Youth	22.9% ± 3.9% 732	20.6% ± 4.3% 754	15.4% ± 2.5% 846	13.4% ± 2.0% 630	20.3% ± 3.7% 1,263

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	33.1% ± 4.9% 797	36.0% ± 6.1% 654	38.1% ± 5.5% 562	31.6% ± 5.3% 448	44.1% ± 7.6% 550
Minority Youth	32.9% ± 12.6% 118	34.3% ± 14.0% 117	41.0% ± 11.2% 156	34.5% ± 11.2% 103	*57.3% ± 11.5% 229

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	8.8% ± 2.3% 699	4.2% ± 1.4% 366	2.6% ± 1.2% 232	2.6% ± 0.5% 230	*2.7% ± 1.0% 237
MS Females	8.2% ± 3.1% 314	3.4% ± 2.0% 148	2.6% ± 1.6% 112	2.2% ± 1.0% 95	*2.5% ± 1.4% 114
MS Males	9.1% ± 2.2% 370	4.8% ± 1.9% 208	2.6% ± 1.5% 120	2.9% ± 0.9% 135	*2.8% ± 1.1% 123
MS Minorities	12.4% ± 5.7% 191	7.1% ± 3.6% 119	3.9% ± 2.7% 88	4.1% ± 1.7% 89	*4.2% ± 1.6% 110
High School (HS)	28.5% ± 3.4% 2,697	22.4% ± 3.1% 2,283	18.8% ± 1.9% 2,239	16.2% ± 2.1% 1,958	*14.0% ± 2.1% 1,692
HS Females	31.3% ± 4.2% 1,448	23.6% ± 3.8% 1,194	18.0% ± 2.2% 1,082	14.9% ± 2.4% 901	*11.6% ± 2.3% 739
HS Males	25.8% ± 3.7% 1,227	21.0% ± 3.8% 1,066	19.4% ± 2.4% 1,146	17.3% ± 2.7% 1,049	*16.6% ± 2.8% 949
HS Minorities	29.3% ± 5.8% 417	27.4% ± 7.4% 473	19.3% ± 3.0% 553	18.0% ± 3.3% 445	*13.7% ± 2.7% 469

Frederick County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Garrett County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	28.1% ± 3.8% 669	27.0% ± 3.9% 639	23.9% ± 3.4% 618	25.0% ± 3.1% 596	28.2% ± 3.4% 628
Any Tobacco Product (Minority Youth)	42.1% ± 13.1% 59	35.9% ± 11.4% 49	32.2% ± 7.8% 69	40.6% ± 8.2% 66	37.3% ± 6.2% 130
Cigarette	22.1% ± 3.1% 513	19.7% ± 3.2% 450	17.0% ± 2.6% 430	17.9% ± 2.7% 416	19.1% ± 2.7% 416
Cigar	9.0% ± 2.0% 212	9.2% ± 2.0% 214	8.4% ± 1.6% 214	11.5% ± 2.0% 273	10.6% ± 1.9% 234
Pipe (tobacco)	4.2% ± 1.4% 99	4.8% ± 1.2% 110	3.5% ± 0.9% 90	6.2% ± 1.5% 146	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.3% ± 1.5% 124	5.6% ± 1.5% 131	3.1% ± 0.8% 80	4.3% ± 1.3% 102	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.2% ± 1.4% 100	4.6% ± 1.2% 107	3.6% ± 0.9% 91	4.2% ± 1.1% 99	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	8.2% ± 2.0% 194	10.0% ± 2.1% 230	9.5% ± 1.8% 242	10.7% ± 2.0% 247	*14.4% ± 2.5% 307

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	24.7% ± 3.2% 585	21.7% ± 2.8% 513	23.5% ± 3.0% 608	21.9% ± 2.7% 522	27.5% ± 3.0% 612
Minority Youth	27.9% ± 13.4% 38	20.2% ± 8.7% 27	23.3% ± 6.0% 50	23.7% ± 6.9% 39	36.0% ± 6.2% 125

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	37.6% ± 7.7% 147	29.6% ± 6.4% 88	33.7% ± 6.5% 82	30.0% ± 5.7% 91	30.4% ± 7.0% 67
Minority Youth	Not Available	Not Available	Not Available	Not Available	35.3% ± 15.5% 15

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.5% ± 3.4% 101	10.9% ± 3.8% 119	8.2% ± 2.6% 93	6.5% ± 2.3% 63	7.1% ± 2.5% 64
MS Females	9.5% ± 4.4% 49	10.0% ± 4.4% 51	10.2% ± 3.7% 55	6.0% ± 3.1% 28	7.5% ± 3.5% 34
MS Males	9.4% ± 5.7% 51	11.6% ± 4.1% 68	6.5% ± 2.9% 39	7.1% ± 2.8% 35	6.8% ± 3.0% 30
MS Minorities	18.4% ± 9.8% 14	11.3% ± 9.6% 8	11.3% ± 5.9% 14	12.9% ± 7.7% 11	11.7% ± 5.5% 20
High School (HS)	32.8% ± 3.8% 413	27.7% ± 4.1% 331	24.1% ± 3.0% 336	25.8% ± 4.2% 353	27.6% ± 3.9% 352
HS Females	31.1% ± 4.9% 184	26.7% ± 5.1% 156	22.6% ± 3.9% 154	25.3% ± 5.3% 173	27.3% ± 5.0% 178
HS Males	34.4% ± 4.9% 228	28.8% ± 5.5% 175	25.5% ± 4.3% 182	26.3% ± 6.0% 181	27.9% ± 5.2% 172
HS Minorities	41.6% ± 15.1% 24	44.6% ± 16.1% 27	36.4% ± 10.9% 26	47.3% ± 10.7% 35	44.3% ± 9.0% 73

Garrett County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Harford County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	26.0% ± 3.3% 5,071	<u>19.7%</u> ± 2.5% 4,056	16.2% ± 2.1% 3,433	15.7% ± 1.8% 3,024	*17.3% ± 1.7% 3,525
Any Tobacco Product (Minority Youth)	25.0% ± 4.5% 1,143	22.5% ± 3.9% 1,184	15.7% ± 3.1% 972	18.8% ± 3.4% 930	18.3% ± 2.3% 1,103
Cigarette	21.2% ± 2.9% 3,933	<u>14.2%</u> ± 2.1% 2,811	11.0% ± 1.7% 2,271	11.0% ± 1.3% 2,079	*10.9% ± 1.3% 2,196
Cigar	11.3% ± 1.9% 2,153	8.3% ± 1.4% 1,661	7.3% ± 1.2% 1,506	9.3% ± 1.4% 1,753	10.0% ± 1.3% 2,021
Pipe (tobacco)	4.7% ± 1.1% 894	4.9% ± 1.1% 997	3.6% ± 0.7% 752	4.8% ± 1.1% 889	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.5% ± 1.2% 1,028	5.1% ± 1.0% 1,031	3.9% ± 0.8% 820	3.4% ± 0.9% 628	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.2% ± 1.0% 808	4.4% ± 1.0% 880	4.4% ± 0.8% 914	3.0% ± 0.7% 562	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.8% ± 1.1% 927	3.7% ± 0.8% 732	3.3% ± 0.7% 681	3.5% ± 0.7% 642	3.3% ± 0.6% 660

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	23.3% ± 2.6% 4,534	<u>17.2%</u> ± 2.1% 3,549	15.8% ± 2.0% 3,343	15.7% ± 1.6% 3,020	<u>19.2%</u> ± 1.7% 3,909
Minority Youth	20.4% ± 3.7% 930	17.5% ± 3.0% 918	12.6% ± 2.1% 781	14.2% ± 2.1% 697	<u>20.0%</u> ± 2.3% 1,206

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	30.0% ± 4.9% 839	31.3% ± 5.4% 521	32.1% ± 5.1% 435	34.3% ± 5.0% 558	40.7% ± 5.9% 549
Minority Youth	34.3% ± 10.2% 195	36.6% ± 13.2% 132	27.8% ± 10.0% 91	38.6% ± 10.1% 172	46.4% ± 9.5% 167

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	10.0% ± 2.6% 874	5.5% ± 1.8% 508	2.7% ± 1.4% 241	3.5% ± 1.2% 307	*2.3% ± 0.8% 201
MS Females	12.7% ± 3.9% 540	5.9% ± 2.4% 271	2.7% ± 1.6% 120	1.9% ± 1.2% 85	*1.4% ± 0.7% 61
MS Males	7.5% ± 2.6% 334	5.1% ± 1.9% 237	2.7% ± 1.6% 121	5.1% ± 1.8% 222	*3.3% ± 1.5% 136
MS Minorities	9.1% ± 4.4% 187	6.3% ± 2.6% 155	3.1% ± 2.4% 85	5.9% ± 2.7% 135	3.3% ± 1.6% 87
High School (HS)	31.0% ± 3.1% 3,059	21.9% ± 3.0% 2,303	17.4% ± 1.9% 2,030	17.3% ± 2.0% 1,772	*17.2% ± 2.0% 1,995
HS Females	31.3% ± 4.2% 1,533	22.9% ± 3.8% 1,216	16.9% ± 2.3% 997	16.5% ± 2.5% 857	*15.7% ± 2.2% 920
HS Males	30.5% ± 3.9% 1,504	20.5% ± 3.2% 1,064	17.8% ± 2.4% 1,033	18.0% ± 2.6% 911	*18.7% ± 2.6% 1,066
HS Minorities	25.7% ± 4.7% 555	21.9% ± 5.0% 542	16.2% ± 3.1% 510	16.1% ± 3.1% 416	*15.1% ± 2.5% 498

Harford County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Howard County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	17.9% ± 3.5% 3,970	17.1% ± 3.5% 4,139	13.3% ± 1.7% 3,603	12.0% ± 1.5% 3,285	12.7% ± 2.6% 3,484
Any Tobacco Product (Minority Youth)	15.7% ± 3.8% 940	17.8% ± 3.5% 1,467	12.4% ± 2.0% 1,426	12.4% ± 1.7% 1,355	14.5% ± 2.8% 1,929
Cigarette	13.5% ± 3.3% 2,913	11.6% ± 2.9% 2,702	8.0% ± 1.2% 2,119	7.8% ± 1.1% 2,094	*7.0% ± 1.7% 1,887
Cigar	7.4% ± 1.8% 1,626	7.4% ± 1.9% 1,749	5.6% ± 0.9% 1,462	7.7% ± 1.2% 2,090	7.0% ± 1.7% 1,911
Pipe (tobacco)	3.6% ± 1.2% 798	4.6% ± 1.2% 1,094	4.1% ± 0.7% 1,101	4.0% ± 0.8% 1,061	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	4.6% ± 1.2% 992	5.3% ± 1.4% 1,252	4.2% ± 0.8% 1,128	<u>2.2%</u> ± 0.5% 602	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.3% ± 1.0% 721	4.7% ± 1.3% 1,112	4.6% ± 0.8% 1,238	<u>2.3%</u> ± 0.5% 614	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.6% ± 1.4% 1,021	4.1% ± 1.2% 972	3.3% ± 0.7% 876	2.5% ± 0.5% 656	*2.3% ± 0.6% 604

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	18.1% ± 2.9% 4,028	15.7% ± 2.9% 3,787	12.9% ± 1.7% 3,497	12.7% ± 1.4% 3,473	14.2% ± 2.6% 3,895
Minority Youth	15.4% ± 3.4% 923	13.4% ± 2.9% 1,107	9.9% ± 1.6% 1,132	10.5% ± 1.7% 1,151	<u>15.9%</u> ± 3.0% 2,110

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	38.5% ± 7.6% 815	34.5% ± 5.9% 594	41.2% ± 6.2% 502	30.8% ± 5.2% 521	<u>46.9%</u> ± 7.8% 473
Minority Youth	39.3% ± 12.1% 194	35.3% ± 10.7% 179	45.1% ± 10.8% 172	31.4% ± 8.1% 216	45.3% ± 10.9% 266

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	4.1% ± 1.1% 407	3.4% ± 1.0% 371	1.7% ± 0.8% 195	1.4% ± 0.8% 168	*1.4% ± 1.0% 155
MS Females	3.2% ± 1.4% 154	3.6% ± 1.0% 193	0.8% ± 0.9% 44	1.4% ± 0.9% 77	1.0% ± 1.2% 59
MS Males	4.9% ± 2.0% 253	3.2% ± 1.3% 178	2.5% ± 1.2% 152	1.5% ± 1.1% 91	*1.8% ± 1.0% 97
MS Minorities	3.6% ± 1.6% 97	5.0% ± 2.6% 186	1.2% ± 1.1% 66	1.6% ± 1.1% 81	2.3% ± 1.7% 133
High School (HS)	21.5% ± 3.0% 2,506	18.7% ± 2.4% 2,331	13.0% ± 1.4% 1,924	12.6% ± 1.8% 1,926	*11.1% ± 1.4% 1,732
HS Females	20.4% ± 3.4% 1,176	18.4% ± 2.9% 1,130	10.4% ± 1.6% 765	10.4% ± 1.9% 788	*9.1% ± 1.6% 757
HS Males	22.5% ± 4.2% 1,310	18.1% ± 2.9% 1,128	15.6% ± 2.0% 1,150	14.5% ± 2.5% 1,123	*13.2% ± 2.1% 962
HS Minorities	18.0% ± 4.6% 540	16.8% ± 3.0% 690	12.2% ± 2.0% 708	11.7% ± 2.3% 686	*12.4% ± 1.8% 900

Howard County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Kent County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	32.0% ± 3.9% 463	29.4% ± 4.0% 424	24.6% ± 4.5% 310	22.8% ± 3.8% 257	23.9% ± 5.0% 254
Any Tobacco Product (Minority Youth)	31.6% ± 5.1% 145	28.0% ± 5.2% 133	22.6% ± 6.5% 101	23.3% ± 5.4% 85	27.9% ± 7.0% 114
Cigarette	23.0% ± 3.8% 317	21.7% ± 3.4% 298	19.1% ± 4.0% 231	17.6% ± 3.1% 195	*14.7% ± 3.9% 153
Cigar	12.4% ± 2.5% 176	11.3% ± 2.1% 158	9.5% ± 2.3% 114	11.9% ± 3.4% 133	12.4% ± 3.2% 130
Pipe (tobacco)	5.8% ± 1.7% 83	8.0% ± 2.0% 112	7.7% ± 2.5% 95	6.2% ± 2.2% 69	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	6.5% ± 1.9% 92	8.2% ± 2.3% 116	7.7% ± 2.4% 96	4.5% ± 2.0% 50	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	5.5% ± 1.5% 78	6.8% ± 2.0% 95	8.1% ± 2.5% 100	4.2% ± 2.0% 47	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	9.3% ± 2.1% 132	8.7% ± 2.0% 121	5.7% ± 1.8% 69	6.9% ± 2.5% 75	7.3% ± 2.1% 73

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	28.7% ± 3.2% 414	24.0% ± 3.1% 348	22.9% ± 3.8% 288	20.6% ± 2.7% 232	23.9% ± 4.3% 254
Minority Youth	28.0% ± 4.6% 128	19.7% ± 4.1% 94	19.2% ± 3.2% 85	17.9% ± 4.8% 65	25.9% ± 5.9% 106

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	36.3% ± 9.3% 84	32.0% ± 7.3% 68	31.2% ± 10.5% 41	29.8% ± 7.8% 46	30.6% ± 10.2% 30
Minority Youth	36.0% ± 13.1% 28	29.1% ± 15.0% 17	Not Available	29.0% ± 13.8% 15	28.7% ± 15.7% 10

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	8.0% ± 2.6% 50	12.9% ± 5.0% 83	6.0% ± 2.7% 28	7.2% ± 3.2% 33	4.1% ± 3.1% 19
MS Females	5.7% ± 3.3% 18	13.6% ± 5.9% 46	3.3% ± 2.7% 8	5.1% ± 3.7% 11	3.4% ± 3.1% 8
MS Males	10.3% ± 3.8% 31	12.1% ± 5.7% 37	8.4% ± 5.3% 20	8.9% ± 4.7% 23	4.2% ± 3.6% 9
MS Minorities	9.3% ± 3.7% 19	12.1% ± 5.4% 25	4.0% ± 3.1% 6	8.0% ± 5.6% 12	*2.7% ± 2.3% 5
High School (HS)	35.3% ± 5.5% 267	29.5% ± 4.1% 215	27.5% ± 3.7% 203	25.2% ± 4.7% 161	*23.1% ± 5.5% 135
HS Females	38.2% ± 6.9% 152	26.5% ± 5.5% 97	24.4% ± 5.9% 87	25.5% ± 5.9% 76	*21.4% ± 6.0% 67
HS Males	32.2% ± 6.6% 115	32.0% ± 5.0% 115	30.6% ± 5.0% 116	24.9% ± 5.3% 86	25.1% ± 7.0% 68
HS Minorities	41.3% ± 7.7% 88	28.8% ± 5.9% 64	22.3% ± 7.2% 58	25.6% ± 7.2% 49	27.6% ± 6.1% 59

Kent County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Montgomery County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	16.6% ± 4.1% 10,604	14.9% ± 2.7% 10,561	14.3% ± 1.7% 10,519	11.5% ± 1.4% 8,348	13.1% ± 2.3% 9,599
Any Tobacco Product (Minority Youth)	15.9% ± 3.4% 4,507	15.0% ± 2.4% 5,755	14.5% ± 1.8% 6,058	11.7% ± 1.6% 4,823	14.5% ± 2.3% 6,730
Cigarette	12.1% ± 3.4% 7,455	9.5% ± 2.1% 6,474	9.0% ± 1.3% 6,414	6.9% ± 0.8% 4,971	*7.1% ± 1.5% 5,182
Cigar	6.6% ± 1.8% 4,107	6.1% ± 1.3% 4,152	6.2% ± 1.0% 4,423	7.7% ± 1.0% 5,529	7.1% ± 1.5% 5,153
Pipe (tobacco)	2.9% ± 0.7% 1,853	3.7% ± 0.8% 2,545	5.0% ± 0.9% 3,640	4.6% ± 0.9% 3,246	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.0% ± 1.6% 3,087	4.6% ± 0.9% 3,177	5.2% ± 0.8% 3,801	<u>3.0%</u> ± 0.5% 2,136	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.4% ± 0.8% 2,113	3.1% ± 0.8% 2,109	<u>5.8%</u> ± 0.9% 4,219	<u>2.5%</u> ± 0.6% 1,755	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	2.9% ± 1.1% 1,836	2.8% ± 0.8% 1,870	3.3% ± 0.7% 2,364	2.6% ± 0.6% 1,849	2.2% ± 0.6% 1,563

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	15.8% ± 3.4% 10,100	13.4% ± 2.8% 9,546	13.1% ± 1.7% 9,640	11.7% ± 1.3% 8,497	15.0% ± 2.5% 11,066
Minority Youth	14.9% ± 2.7% 4,232	12.4% ± 2.5% 4,785	11.8% ± 1.6% 4,906	10.6% ± 1.1% 4,355	<u>16.3%</u> ± 2.4% 7,581

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	36.9% ± 6.9% 1,955	43.7% ± 3.5% 2,046	45.0% ± 6.2% 1,577	39.4% ± 5.6% 1,481	*50.8% ± 6.4% 1,881
Minority Youth	46.7% ± 12.1% 987	49.4% ± 5.3% 1,218	47.9% ± 7.9% 995	41.6% ± 6.1% 1,010	52.3% ± 6.2% 1,452

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	3.7% ± 1.5% 1,065	3.1% ± 1.0% 981	3.0% ± 1.2% 916	1.3% ± 0.7% 408	2.7% ± 1.2% 816
MS Females	3.3% ± 2.1% 450	2.3% ± 0.9% 346	2.1% ± 1.3% 320	1.0% ± 0.7% 146	2.3% ± 1.6% 362
MS Males	3.8% ± 2.5% 558	4.0% ± 1.8% 635	3.6% ± 1.8% 556	1.7% ± 1.1% 262	3.1% ± 1.5% 454
MS Minorities	4.4% ± 1.7% 566	3.9% ± 1.2% 685	3.3% ± 1.5% 553	1.8% ± 0.9% 316	3.7% ± 1.7% 706
High School (HS)	19.4% ± 2.5% 6,389	14.9% ± 2.0% 5,494	13.4% ± 1.5% 5,498	11.1% ± 1.3% 4,563	*10.4% ± 1.5% 4,366
HS Females	19.6% ± 4.4% 3,217	14.1% ± 2.5% 2,583	12.8% ± 2.0% 2,573	8.9% ± 1.3% 1,823	*8.2% ± 1.7% 1,798
HS Males	18.8% ± 3.9% 3,101	15.4% ± 1.9% 2,836	14.0% ± 1.9% 2,899	13.1% ± 2.0% 2,700	*12.7% ± 1.9% 2,568
HS Minorities	15.4% ± 2.9% 2,222	14.0% ± 1.9% 2,683	12.9% ± 1.7% 3,006	11.2% ± 1.3% 2,543	*11.1% ± 1.3% 2,927

Montgomery County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Prince George's County Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	17.1% ± 3.2% 10,459	15.3% ± 2.5% 10,097	11.9% ± 1.5% 8,341	<u>15.3%</u> ± 1.5% 10,089	17.4% ± 2.2% 11,020
Any Tobacco Product (Minority Youth)	15.7% ± 3.2% 8,126	14.5% ± 2.3% 8,236	11.3% ± 1.4% 7,347	<u>14.8%</u> ± 1.5% 9,105	17.2% ± 2.1% 10,419
Cigarette	10.6% ± 2.5% 6,178	8.5% ± 1.7% 5,269	6.3% ± 1.0% 4,254	7.7% ± 0.9% 4,964	7.6% ± 1.2% 4,715
Cigar	7.4% ± 1.7% 4,369	6.1% ± 1.4% 3,784	4.4% ± 0.8% 2,933	<u>11.2%</u> ± 1.4% 7,170	9.9% ± 1.6% 6,191
Pipe (tobacco)	4.2% ± 1.5% 2,487	4.8% ± 1.2% 2,973	4.5% ± 0.9% 3,100	6.7% ± 1.4% 4,159	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.8% ± 1.4% 3,357	5.0% ± 1.2% 3,174	4.6% ± 0.9% 3,149	4.9% ± 1.0% 2,998	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	2.9% ± 1.1% 1,743	4.2% ± 1.1% 2,592	5.1% ± 1.0% 3,467	4.4% ± 1.1% 2,691	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	2.4% ± 1.0% 1,423	2.9% ± 0.9% 1,808	2.6% ± 0.6% 1,759	3.2% ± 0.6% 1,947	2.8% ± 0.6% 1,631

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	14.9% ± 2.0% 9,115	13.3% ± 1.7% 8,759	10.8% ± 1.3% 7,547	12.7% ± 0.8% 8,371	<u>17.9%</u> ± 1.9% 11,346
Minority Youth	14.0% ± 2.1% 7,262	12.6% ± 1.6% 7,149	10.4% ± 1.3% 6,789	12.3% ± 1.0% 7,545	<u>17.7%</u> ± 1.9% 10,757

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	39.1% ± 7.9% 1,839	43.9% ± 6.4% 1,686	51.6% ± 7.0% 1,840	52.0% ± 6.0% 2,974	*55.3% ± 4.9% 2,171
Minority Youth	48.9% ± 11.2% 1,639	47.5% ± 6.6% 1,431	53.9% ± 7.5% 1,731	55.1% ± 6.4% 2,867	55.0% ± 5.2% 2,064

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	5.1% ± 3.0% 1,363	3.8% ± 1.3% 1,098	3.7% ± 1.4% 1,063	3.4% ± 1.3% 925	3.9% ± 1.2% 1,001
MS Females	3.1% ± 1.0% 411	3.7% ± 2.3% 536	3.4% ± 2.3% 486	2.2% ± 0.9% 300	4.1% ± 1.2% 548
MS Males	6.9% ± 4.9% 925	3.7% ± 0.9% 541	3.9% ± 1.7% 577	4.6% ± 2.5% 625	3.6% ± 1.7% 453
MS Minorities	5.0% ± 3.4% 1,136	3.7% ± 1.4% 910	3.8% ± 1.5% 1,019	3.0% ± 1.3% 761	3.8% ± 1.3% 944
High School (HS)	15.3% ± 3.0% 4,816	12.8% ± 1.5% 4,172	8.3% ± 1.1% 3,191	10.9% ± 1.2% 4,039	*10.2% ± 1.2% 3,714
HS Females	16.0% ± 3.6% 2,577	12.1% ± 2.2% 2,033	6.4% ± 1.3% 1,250	8.0% ± 1.4% 1,475	*6.7% ± 1.1% 1,268
HS Males	14.1% ± 3.4% 2,155	12.8% ± 2.2% 2,011	10.3% ± 1.6% 1,941	13.9% ± 1.8% 2,558	14.0% ± 2.1% 2,423
HS Minorities	12.1% ± 3.0% 3,202	11.0% ± 1.4% 3,121	7.4% ± 1.0% 2,615	10.3% ± 1.4% 3,545	9.9% ± 1.3% 3,419

Prince George's County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Queen Anne's County Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	25.7% ± 3.7% 896	22.4% ± 2.2% 833	21.4% ± 2.8% 899	19.1% ± 2.1% 802	20.5% ± 2.0% 819
Any Tobacco Product (Minority Youth)	32.3% ± 6.2% 170	30.1% ± 4.8% 172	31.3% ± 4.9% 207	32.7% ± 5.1% 220	30.2% ± 4.2% 265
Cigarette	20.4% ± 3.3% 683	17.4% ± 2.0% 622	15.7% ± 2.4% 641	14.7% ± 1.8% 610	*13.9% ± 1.7% 547
Cigar	10.0% ± 1.8% 342	7.8% ± 1.2% 283	9.4% ± 1.5% 381	10.8% ± 1.6% 450	10.7% ± 1.5% 426
Pipe (tobacco)	4.3% ± 1.1% 148	5.3% ± 1.2% 194	5.2% ± 1.1% 217	7.5% ± 1.3% 311	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	4.6% ± 1.1% 157	6.3% ± 1.2% 230	5.7% ± 1.1% 236	5.0% ± 1.0% 208	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.8% ± 1.0% 132	5.8% ± 1.3% 209	5.9% ± 1.2% 243	5.2% ± 1.0% 214	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.3% ± 1.0% 149	5.3% ± 1.1% 192	5.0% ± 1.0% 204	5.2% ± 0.9% 208	5.0% ± 0.9% 191

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	25.0% ± 3.4% 872	<u>18.1%</u> ± 1.7% 675	20.3% ± 2.5% 850	16.3% ± 1.6% 685	<u>21.2%</u> ± 1.9% 847
Minority Youth	26.7% ± 7.3% 141	19.4% ± 3.9% 111	20.6% ± 3.7% 135	18.4% ± 3.3% 123	<u>29.8%</u> ± 3.6% 261

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	24.8% ± 5.3% 115	28.2% ± 5.4% 109	29.5% ± 5.8% 114	28.0% ± 5.3% 113	*38.8% ± 5.8% 114
Minority Youth	26.5% ± 11.0% 19	27.3% ± 12.2% 13	25.9% ± 11.5% 16	28.3% ± 11.8% 19	31.1% ± 12.8% 20

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.6% ± 3.6% 152	5.3% ± 1.8% 91	4.3% ± 1.3% 74	2.8% ± 1.0% 49	*3.1% ± 1.0% 52
MS Females	8.3% ± 3.7% 62	5.4% ± 2.4% 44	4.1% ± 1.6% 34	2.3% ± 0.9% 19	*2.2% ± 1.1% 18
MS Males	10.8% ± 4.5% 91	5.2% ± 10.9% 47	4.6% ± 1.7% 40	3.3% ± 1.6% 30	*4.1% ± 1.5% 34
MS Minorities	15.3% ± 7.9% 35	7.9% ± 3.3% 20	11.0% ± 4.2% 27	5.3% ± 2.7% 14	5.2% ± 2.5% 18
High School (HS)	30.1% ± 3.1% 531	28.6% ± 3.1% 531	23.8% ± 3.0% 567	23.5% ± 2.9% 561	*21.8% ± 2.7% 495
HS Females	30.9% ± 4.5% 261	29.1% ± 3.8% 260	23.6% ± 4.0% 289	20.7% ± 3.4% 245	*19.3% ± 3.1% 227
HS Males	29.5% ± 4.2% 268	27.8% ± 4.6% 263	24.1% ± 3.2% 278	26.3% ± 3.7% 317	24.1% ± 3.7% 261
HS Minorities	33.5% ± 6.9% 85	33.0% ± 7.4% 92	27.8% ± 5.7% 101	35.2% ± 7.0% 134	33.5% ± 5.6% 161

Queen Anne's County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Somerset County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	33.9% ± 4.1% 504	27.3% ± 4.3% 390	20.4% ± 3.4% 303	23.2% ± 3.2% 326	*25.5% ± 3.9% 331
Any Tobacco Product (Minority Youth)	30.6% ± 4.4% 202	27.3% ± 5.2% 190	<u>15.1%</u> ± 3.5% 114	<u>24.6%</u> ± 4.7% 179	24.9% ± 3.8% 178
Cigarette	29.2% ± 4.5% 415	<u>20.7%</u> ± 3.7% 275	<u>13.5%</u> ± 3.1% 194	15.5% ± 2.4% 213	*17.3% ± 3.2% 219
Cigar	14.6% ± 3.4% 210	11.2% ± 2.5% 152	8.5% ± 1.9% 120	<u>16.6%</u> ± 2.9% 228	14.7% ± 3.0% 188
Pipe (tobacco)	6.4% ± 2.1% 94	8.3% ± 2.3% 113	5.7% ± 1.5% 82	<u>9.9%</u> ± 2.5% 135	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	6.4% ± 1.7% 91	7.6% ± 2.0% 103	5.3% ± 1.5% 76	7.0% ± 1.6% 95	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	5.6% ± 1.9% 82	7.3% ± 2.3% 99	6.1% ± 1.7% 89	5.7% ± 1.7% 78	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	5.4% ± 1.8% 78	6.5% ± 2.8% 86	4.6% ± 1.4% 65	5.2% ± 1.5% 68	6.5% ± 2.0% 79

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	26.4% ± 3.1% 393	23.3% ± 3.4% 334	18.3% ± 2.5% 272	21.4% ± 3.5% 299	26.4% ± 3.8% 343
Minority Youth	24.6% ± 3.5% 162	22.1% ± 4.1% 155	<u>12.8%</u> ± 2.6% 97	<u>23.0%</u> ± 4.3% 166	25.6% ± 4.3% 183

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	38.7% ± 7.8% 121	39.4% ± 8.4% 67	28.1% ± 9.6% 36	37.6% ± 7.6% 64	37.2% ± 7.5% 56
Minority Youth	32.8% ± 10.8% 30	44.4% ± 18.1% 25	Not Available	42.8% ± 11.9% 37	41.5% ± 10.8% 31

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	17.5% ± 4.3% 113	14.4% ± 4.9% 88	6.9% ± 2.3% 47	8.3% ± 2.9% 53	*7.6% ± 2.9% 42
MS Females	13.1% ± 6.9% 41	10.0% ± 5.2% 31	8.9% ± 3.6% 29	6.9% ± 3.2% 20	5.0% ± 2.5% 14
MS Males	21.1% ± 5.1% 68	18.6% ± 6.2% 54	5.2% ± 2.0% 19	9.3% ± 4.5% 32	*10.2% ± 4.5% 28
MS Minorities	16.0% ± 6.1% 48	12.1% ± 5.6% 38	5.3% ± 2.5% 18	11.6% ± 4.6% 38	7.1% ± 3.2% 23
High School (HS)	38.9% ± 8.0% 302	26.1% ± 5.4% 187	19.5% ± 4.4% 147	21.5% ± 3.4% 160	*24.9% ± 5.0% 177
HS Females	35.1% ± 10.3% 142	21.4% ± 6.0% 83	17.0% ± 4.7% 69	14.7% ± 4.1% 55	25.0% ± 5.3% 93
HS Males	42.5% ± 8.0% 158	31.2% ± 7.2% 101	22.3% ± 6.6% 78	28.3% ± 5.9% 104	*24.9% ± 6.5% 84
HS Minorities	37.2% ± 7.8% 117	22.3% ± 6.9% 70	10.7% ± 4.7% 41	16.6% ± 5.0% 63	*22.7% ± 4.8% 83

Somerset County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





St. Mary's County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	24.0% ± 3.0% 1,776	23.7% ± 3.2% 1,841	<u>14.7%</u> ± 1.9% 1,282	16.1% ± 2.0% 1,402	*15.7% ± 1.9% 1,369
Any Tobacco Product (Minority Youth)	22.6% ± 4.0% 483	26.1% ± 5.3% 583	<u>16.3%</u> ± 3.0% 430	18.1% ± 3.1% 418	18.2% ± 3.0% 458
Cigarette	19.4% ± 2.9% 1,363	18.0% ± 2.9% 1,345	<u>10.2%</u> ± 1.6% 870	11.8% ± 1.6% 1,012	*10.1% ± 1.4% 864
Cigar	10.1% ± 1.6% 729	8.0% ± 1.6% 603	5.8% ± 1.0% 490	<u>9.5%</u> ± 1.5% 813	8.1% ± 1.4% 696
Pipe (tobacco)	3.9% ± 1.0% 286	5.0% ± 1.6% 377	3.5% ± 0.7% 299	5.2% ± 1.0% 444	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	4.9% ± 1.2% 345	5.6% ± 1.5% 420	3.3% ± 0.7% 286	3.7% ± 0.8% 316	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.2% ± 1.0% 234	5.3% ± 1.6% 399	3.7% ± 0.7% 314	3.7% ± 0.8% 311	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.5% ± 1.0% 255	4.2% ± 1.5% 309	2.7% ± 0.6% 234	3.5% ± 0.8% 296	3.3% ± 0.7% 274

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	22.5% ± 2.6% 1,662	19.4% ± 2.0% 1,511	15.5% ± 2.0% 1,346	16.3% ± 2.0% 1,421	*16.5% ± 1.9% 1,435
Minority Youth	23.1% ± 4.3% 492	19.6% ± 4.2% 438	14.8% ± 2.9% 390	15.4% ± 2.6% 354	19.4% ± 3.0% 488

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	32.7% ± 6.0% 322	25.2% ± 5.3% 202	<u>39.2%</u> ± 6.4% 247	35.2% ± 5.5% 281	*45.0% ± 6.2% 253
Minority Youth	37.2% ± 9.6% 66	<u>15.7%</u> ± 10.3% 26	<u>40.4%</u> ± 12.9% 62	40.1% ± 13.4% 79	51.3% ± 10.7% 89

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	7.5% ± 2.5% 235	8.5% ± 2.1% 285	<u>3.2%</u> ± 1.3% 114	4.3% ± 1.8% 153	*3.7% ± 1.2% 139
MS Females	6.1% ± 2.8% 91	7.5% ± 2.8% 127	<u>2.7%</u> ± 1.5% 48	3.2% ± 2.1% 56	3.4% ± 1.4% 64
MS Males	8.6% ± 3.1% 141	9.2% ± 2.9% 154	<u>3.7%</u> ± 1.8% 66	5.3% ± 2.3% 97	4.1% ± 1.8% 76
MS Minorities	9.3% ± 4.2% 92	9.9% ± 3.6% 99	4.1% ± 2.4% 46	5.1% ± 2.6% 48	3.1% ± 1.8% 33
High School (HS)	29.0% ± 3.8% 1,129	26.0% ± 4.5% 1,060	<u>15.4%</u> ± 1.9% 756	17.2% ± 2.4% 859	*14.9% ± 2.1% 724
HS Females	26.8% ± 3.9% 524	30.4% ± 6.7% 624	<u>15.5%</u> ± 2.2% 387	15.9% ± 2.9% 389	*12.6% ± 2.5% 313
HS Males	31.1% ± 5.1% 596	21.6% ± 5.5% 433	15.4% ± 2.5% 369	18.5% ± 3.2% 468	*17.2% ± 2.7% 406
HS Minorities	20.4% ± 4.7% 201	22.9% ± 7.4% 249	14.1% ± 2.7% 199	16.7% ± 4.0% 220	15.4% ± 3.6% 213

St. Mary's County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Talbot County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	28.3% ± 3.2% 633	26.1% ± 3.1% 606	23.5% ± 3.5% 570	23.0% ± 2.9% 547	*21.4% ± 3.2% 483
Any Tobacco Product (Minority Youth)	24.1% ± 4.3% 146	25.6% ± 4.6% 170	22.6% ± 5.2% 171	26.7% ± 4.8% 198	25.2% ± 4.8% 196
Cigarette	23.1% ± 3.1% 494	18.1% ± 2.4% 399	17.7% ± 3.0% 411	17.3% ± 2.5% 400	*15.0% ± 2.4% 331
Cigar	12.6% ± 2.0% 276	12.1% ± 2.1% 270	8.6% ± 1.6% 200	13.3% ± 2.3% 310	10.6% ± 2.1% 237
Pipe (tobacco)	5.7% ± 1.5% 127	6.9% ± 1.8% 155	6.7% ± 1.6% 160	8.4% ± 1.8% 195	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.6% ± 1.3% 120	7.7% ± 1.9% 173	6.9% ± 1.9% 164	5.8% ± 1.5% 134	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.5% ± 1.2% 98	6.6% ± 1.6% 147	7.2% ± 1.8% 172	5.6% ± 1.2% 131	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.1% ± 1.1% 90	5.7% ± 1.2% 125	6.0% ± 1.4% 140	6.5% ± 1.5% 146	6.1% ± 1.4% 131

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	26.7% ± 3.2% 598	<u>20.7%</u> ± 2.3% 480	21.7% ± 3.2% 525	19.3% ± 2.1% 457	21.6% ± 2.7% 487
Minority Youth	23.5% ± 5.0% 142	17.9% ± 3.3% 118	15.5% ± 3.8% 116	17.7% ± 3.6% 131	23.7% ± 4.1% 184

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	26.5% ± 6.0% 84	33.6% ± 7.0% 74	35.1% ± 6.3% 100	29.5% ± 6.3% 78	*42.0% ± 9.3% 68
Minority Youth	38.2% ± 11.8% 27	26.8% ± 14.7% 10	44.6% ± 14.7% 28	32.7% ± 11.5% 28	34.8% ± 15.3% 17

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*’.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.3% ± 4.1% 91	7.2% ± 2.7% 72	5.9% ± 1.9% 57	7.5% ± 3.1% 70	4.0% ± 1.9% 36
MS Females	8.7% ± 4.9% 41	6.4% ± 2.6% 30	4.4% ± 1.9% 22	6.9% ± 3.7% 33	3.2% ± 2.2% 14
MS Males	10.0% ± 4.4% 50	8.1% ± 3.3% 42	7.5% ± 2.8% 35	8.1% ± 3.8% 38	4.8% ± 2.6% 22
MS Minorities	11.0% ± 6.2% 31	8.1% ± 4.2% 26	6.1% ± 4.0% 19	9.5% ± 5.4% 29	3.6% ± 2.4% 11
High School (HS)	34.5% ± 3.8% 403	27.0% ± 3.4% 327	26.2% ± 3.4% 354	23.9% ± 3.6% 330	*22.4% ± 3.4% 296
HS Females	32.5% ± 5.1% 177	26.1% ± 4.7% 153	25.2% ± 4.7% 160	23.2% ± 5.3% 160	*19.4% ± 3.6% 129
HS Males	36.1% ± 4.6% 224	27.5% ± 4.0% 170	26.8% ± 4.3% 191	24.5% ± 3.7% 169	*25.5% ± 5.4% 165
HS Minorities	22.5% ± 5.0% 61	26.4% ± 5.4% 77	19.9% ± 6.1% 77	26.1% ± 5.8% 106	25.8% ± 5.3% 116

Talbot County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Washington County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	28.0% ± 3.5% 2,770	22.4% ± 2.7% 2,321	20.9% ± 2.5% 2,319	20.0% ± 2.0% 2,152	*21.7% ± 2.5% 2,457
Any Tobacco Product (Minority Youth)	29.5% ± 6.6% 446	28.4% ± 4.5% 464	21.8% ± 3.4% 532	20.4% ± 3.7% 421	27.3% ± 4.1% 730
Cigarette	22.9% ± 3.2% 2,184	17.5% ± 2.6% 1,754	14.9% ± 2.1% 1,609	14.5% ± 1.8% 1,535	*14.2% ± 2.0% 1,580
Cigar	9.9% ± 1.6% 967	7.2% ± 1.3% 735	8.5% ± 1.3% 923	11.2% ± 1.4% 1,182	10.5% ± 1.5% 1,165
Pipe (tobacco)	3.6% ± 0.8% 358	4.0% ± 0.9% 408	4.7% ± 0.9% 518	6.0% ± 1.1% 620	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	4.4% ± 1.1% 427	4.3% ± 0.9% 438	4.4% ± 0.8% 482	3.8% ± 0.8% 389	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.7% ± 0.9% 364	3.4% ± 0.9% 348	4.6% ± 0.8% 508	4.0% ± 0.9% 415	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	5.8% ± 1.3% 563	4.6% ± 1.1% 461	5.1% ± 1.0% 553	6.2% ± 0.9% 624	6.7% ± 1.2% 709

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	26.0% ± 2.9% 2,569	22.9% ± 2.5% 2,373	22.3% ± 2.5% 2,484	19.1% ± 1.8% 2,054	21.1% ± 2.3% 2,387
Minority Youth	27.6% ± 5.5% 417	24.3% ± 5.0% 398	20.3% ± 3.5% 495	17.3% ± 3.3% 359	<u>25.6%</u> ± 3.7% 685

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	36.6% ± 4.5% 622	40.2% ± 5.9% 517	37.6% ± 4.8% 434	34.3% ± 5.1% 435	35.2% ± 5.5% 382
Minority Youth	47.2% ± 14.9% 120	36.4% ± 14.2% 74	44.2% ± 12.1% 94	34.2% ± 11.0% 83	40.4% ± 9.7% 105

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '*'.


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	12.6% ± 3.9% 559	8.9% ± 2.7% 408	5.4% ± 1.5% 253	4.1% ± 1.6% 197	*4.9% ± 1.7% 242
MS Females	12.0% ± 4.8% 259	8.1% ± 3.0% 183	4.9% ± 2.1% 110	3.2% ± 1.5% 77	*3.6% ± 1.8% 91
MS Males	13.2% ± 4.5% 301	9.6% ± 3.4% 222	5.9% ± 1.9% 143	4.9% ± 2.3% 120	6.0% ± 2.5% 144
MS Minorities	10.9% ± 7.4% 83	16.7% ± 5.7% 142	6.8% ± 3.0% 80	4.9% ± 2.8% 50	6.3% ± 3.2% 76
High School (HS)	31.9% ± 3.1% 1,624	24.8% ± 3.3% 1,345	22.1% ± 2.3% 1,356	23.2% ± 2.8% 1,339	*21.8% ± 2.6% 1,337
HS Females	34.3% ± 4.3% 862	26.2% ± 4.4% 717	23.0% ± 2.7% 699	23.0% ± 3.3% 646	*23.6% ± 3.6% 757
HS Males	29.6% ± 3.7% 758	23.3% ± 4.0% 617	21.1% ± 3.0% 654	23.3% ± 3.4% 690	*19.7% ± 2.7% 576
HS Minorities	31.4% ± 6.5% 205	26.1% ± 6.4% 172	20.8% ± 3.7% 234	24.1% ± 4.6% 245	25.3% ± 3.9% 349

Washington County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Wicomico County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	27.0% ± 3.3% 1,817	23.1% ± 3.5% 1,554	<u>17.1%</u> ± 2.0% 1,196	19.0% ± 1.8% 1,298	22.5% ± 2.6% 1,514
Any Tobacco Product (Minority Youth)	29.1% ± 4.5% 673	21.4% ± 4.0% 599	16.8% ± 2.5% 555	18.7% ± 2.2% 572	<u>24.5%</u> ± 3.0% 801
Cigarette	22.4% ± 3.1% 1,450	18.8% ± 3.1% 1,214	<u>12.0%</u> ± 1.6% 814	12.9% ± 1.3% 865	*15.2% ± 2.0% 997
Cigar	12.7% ± 2.0% 835	8.9% ± 2.1% 577	7.9% ± 1.3% 532	<u>12.4%</u> ± 1.5% 823	13.5% ± 1.9% 891
Pipe (tobacco)	4.0% ± 1.1% 267	5.4% ± 1.7% 350	3.6% ± 0.8% 247	<u>5.9%</u> ± 1.1% 386	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	3.9% ± 1.1% 256	4.3% ± 1.4% 281	3.6% ± 0.9% 243	4.0% ± 0.8% 262	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	3.4% ± 1.1% 229	3.3% ± 1.2% 212	4.1% ± 1.0% 276	3.5% ± 0.9% 229	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	2.5% ± 0.8% 169	3.3% ± 1.0% 212	3.3% ± 0.7% 225	4.0% ± 0.8% 258	*5.1% ± 1.0% 316

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	24.6% ± 2.5% 1,656	20.9% ± 3.2% 1,407	18.9% ± 2.1% 1,332	18.6% ± 1.8% 1,266	22.3% ± 2.4% 1,497
Minority Youth	23.8% ± 3.0% 552	<u>16.8%</u> ± 3.8% 471	15.8% ± 2.1% 526	17.1% ± 2.1% 519	<u>23.4%</u> ± 2.8% 768

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	34.3% ± 5.7% 370	35.0% ± 8.2% 284	40.6% ± 5.9% 240	36.7% ± 5.9% 248	33.1% ± 5.5% 187
Minority Youth	31.0% ± 8.0% 116	36.1% ± 13.6% 94	41.7% ± 10.4% 91	44.7% ± 8.8% 128	36.0% ± 7.8% 93

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	12.0% ± 3.7% 360	10.4% ± 3.2% 299	6.4% ± 1.7% 184	6.4% ± 1.6% 183	7.2% ± 2.6% 212
MS Females	12.9% ± 4.7% 173	11.0% ± 4.4% 153	5.5% ± 2.4% 77	5.8% ± 2.0% 79	5.7% ± 2.5% 86
MS Males	11.3% ± 4.1% 184	9.8% ± 3.6% 147	7.2% ± 2.7% 108	7.0% ± 2.7% 104	8.7% ± 3.4% 125
MS Minorities	16.8% ± 5.4% 183	8.6% ± 3.0% 116	6.8% ± 2.4% 97	6.6% ± 2.4% 91	8.8% ± 3.5% 130
High School (HS)	31.3% ± 3.9% 1,090	25.6% ± 4.4% 914	16.2% ± 1.9% 629	17.8% ± 1.9% 682	*21.6% ± 2.6% 785
HS Females	32.4% ± 5.1% 572	24.4% ± 5.7% 418	16.5% ± 2.5% 327	16.7% ± 2.5% 322	*19.3% ± 3.3% 369
HS Males	30.1% ± 5.0% 512	26.1% ± 4.9% 477	15.9% ± 2.6% 302	18.9% ± 2.6% 359	24.2% ± 3.0% 414
HS Minorities	29.6% ± 4.6% 332	21.2% ± 5.8% 279	13.7% ± 2.8% 234	13.9% ± 2.4% 221	*20.5% ± 2.7% 352

Wicomico County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Worcester County

Tobacco Use by Youth: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Youth Populations)	25.2% ± 3.5% 901	23.0% ± 3.1% 849	19.3% ± 2.4% 729	22.5% ± 1.9% 784	23.9% ± 2.5% 840
Any Tobacco Product (Minority Youth)	23.1% ± 4.5% 238	30.1% ± 6.0% 339	<u>19.7%</u> ± 3.1% 240	24.6% ± 3.2% 258	26.2% ± 3.1% 310
Cigarette	19.9% ± 3.1% 683	16.3% ± 3.0% 575	13.7% ± 2.0% 505	15.6% ± 1.6% 533	16.1% ± 2.1% 551
Cigar	10.2% ± 2.1% 356	11.0% ± 2.2% 395	8.9% ± 1.4% 328	<u>13.9%</u> ± 1.4% 484	12.8% ± 1.9% 449
Pipe (tobacco)	4.7% ± 1.2% 164	6.4% ± 1.6% 228	4.4% ± 0.8% 166	5.4% ± 1.1% 186	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010
Bidi	5.2% ± 1.3% 180	5.6% ± 1.9% 201	3.9% ± 0.8% 145	3.7% ± 1.0% 129	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010
Kretek	4.1% ± 1.2% 143	5.4% ± 2.3% 193	4.0% ± 0.8% 148	3.4% ± 0.8% 118	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.7% ± 1.1% 131	5.0% ± 1.5% 176	4.3% ± 0.9% 161	5.3% ± 0.9% 181	*6.7% ± 1.2% 229

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	21.4% ± 2.7% 765	19.2% ± 2.5% 710	19.7% ± 2.4% 743	22.5% ± 1.9% 785	23.3% ± 2.2% 817
Minority Youth	17.6% ± 3.7% 181	18.8% ± 4.3% 211	17.3% ± 3.0% 212	19.8% ± 2.9% 207	*24.6% ± 2.9% 291

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
All Youth	34.3% ± 5.7% 370	35.0% ± 8.2% 284	40.6% ± 5.9% 240	36.7% ± 5.9% 248	38.1% ± 6.4% 149
Minority Youth	42.7% ± 14.3% 43	28.9% ± 11.8% 28	38.2% ± 13.0% 30	35.2% ± 9.6% 38	43.7% ± 9.3% 53

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2010 preceded by an asterisk '**'


CURRENT CIGARETTE SMOKING — School Type, Gender & Minorities

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010
Middle School (MS)	9.0% ± 3.2% 133	6.4% ± 1.4% 100	5.0% ± 1.4% 71	4.2% ± 1.2% 58	5.8% ± 1.8% 82
MS Females	5.7% ± 2.9% 40	6.0% ± 1.9% 47	5.3% ± 1.7% 38	3.4% ± 1.4% 23	6.6% ± 2.3% 48
MS Males	12.2% ± 4.4% 93	6.9% ± 2.2% 53	4.7% ± 1.7% 33	4.9% ± 1.8% 34	*4.7% ± 2.0% 33
MS Minorities	8.1% ± 4.1% 32	7.8% ± 2.3% 36	6.3% ± 2.5% 30	3.9% ± 2.0% 16	7.6% ± 2.9% 39
High School (HS)	28.0% ± 3.9% 550	24.0% ± 4.6% 475	19.2% ± 2.4% 433	23.3% ± 2.5% 475	23.4% ± 3.1% 468
HS Females	24.6% ± 5.2% 229	23.0% ± 6.2% 219	18.8% ± 2.7% 208	20.3% ± 3.2% 204	19.6% ± 3.4% 205
HS Males	31.1% ± 4.8% 320	24.5% ± 5.6% 249	19.6% ± 2.6% 226	26.3% ± 3.3% 271	27.4% ± 5.3% 264
HS Minorities	20.5% ± 5.9% 118	28.3% ± 8.1% 170	16.9% ± 3.9% 117	21.1% ± 3.8% 125	21.5% ± 3.6% 135

Worcester County - Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)



Appendix B

Adult Prevalence, Initiation, and Cessation

	<u>Page</u>
Maryland	88
Allegany	90
Anne Arundel	92
Baltimore City	94
Baltimore County	96
Calvert	98
Caroline	100
Carroll	102
Cecil	104
Charles	106
Dorchester	108
Frederick	110
Garrett	112
Harford	114
Howard	116
Kent	118
Montgomery	120
Prince George's	122
Queen Anne's	124
Somerset	126
St. Mary's	128
Talbot	130
Washington	132
Wicomico	134
Worcester	136



Maryland

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	20.5% ± 1.5% 799,273	21.9% ± 1.7% 879,623	<u>17.7%</u> ± 1.2% 746,347	<u>14.9%</u> ± 1.1% 641,382	*15.2% ± 1.2% 654,019
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.4% ± 0.4% 54,102	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.1% ± 0.5% 90,581

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	6.2% ± 2.9% 10,727	7.6% ± 3.3% 11,350	9.5% ± 4.5% 15,267	7.0% ± 3.8% 8,557	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	23.4% ± 3.4% 73,606	22.2% ± 3.7% 67,716	<u>31.7%</u> ± 4.1% 88,244	29.7% ± 3.5% 95,037	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Selected Populations

Proportion (%) of Selected Populations and Number For Pregnant Women


Populations Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Pregnant Women ⁴ 100% of Maryland Birth Certificates	9.2% 6,842	8.0% 5,877	6.8% 5,261	6.6% 5,105	6.1% 4,584
Black/African American ⁵	21.7% ± 3.4% 213,250	23.8% ± 4.3% 246,663	19.8% ± 3.0% 214,319	16.4% ± 2.6% 188,482	17.8% ± 2.7% 215,562
Hispanic ⁵	17.2% ± 7.8% 31,948	17.8% ± 10.4% 34,346	11.1% ± 5.9% 22,190	7.9% ± 4.2% 21,759	7.8% ± 4.9% 18,562
White ⁵	21.0% ± 1.7% 528,634	21.4% ± 1.7% 533,766	17.9% ± 1.3% 448,145	15.8% ± 1.3% 395,639	*15.0% ± 1.4% 373,246

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁶

Current Adult (Age 18 and older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	37.9% ± 10.9% 10,495	40.3% ± 9.2% 7,434	37.8% ± 9.4% 7,158	44.6% ± 9.0% 8,278	DATA NOT AVAIL- ABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁷ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

7 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Allegany County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	26.8% ± 9.0% 13,738	24.1% ± 8.4% 15,237	27.5% ± 9.3% 16,729	23.5% ± 7.0% 14,606	20.9% ± 9.0% 14,066
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	5.8% ± 4.5% 2,936	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	6.0% ± 4.9% 4,028

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	1.4% ± 2.1% 65	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	34.1% ± 23.7% 1,872	26.9% ± 16.5% 1,651	37.8% ± 23.8% 2,065	22.5% ± 13.4% 1,405	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Allegany County	16.8% 133	18.1% 128	22.1% 157	22.7% 153	24.8% 159	20.2% 131	23.7% 166	19.0% 130	24.7% 176	24.1% 178

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	37.9% ± 10.9% 10,495	40.3% ± 9.2% 7,434	37.8% ± 9.4% 7,158	44.6% ± 9.0% 8,278	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Anne Arundel County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	25.0% ± 5.7% 87,584	23.8% ± 5.6% 90,944	16.9% ± 3.9% 60,077	13.8% ± 4.1% 52,114	15.4% ± 4.4% 57,566
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.3% ± 1.9% 4,449	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	1.7% ± 1.4% 6,243

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	19.1% ± 16.4% 3,317	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	28.0% ± 10.2% 9,137	18.1% ± 9.9% 4,810	28.9% ± 11.6% 7,071	24.9% ± 10.4% 6,592	DATA NOT AVAILABLE FOR 2010

1 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

2 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

3 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Anne Arundel County	11.1% 754	10.5% 724	9.7% 656	8.5% 589	8.7% 587	7.6% 514	8.1% 576	7.7% 547	7.7% 552	6.7% 478

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	37.3% ± 5.3% 54,805	31.8% ± 5.4% 47,214	27.8% ± 4.5% 42,400	23.2% ± 4.2% 36,921	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Baltimore City

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	29.7% ± 5.9% 131,434	25.3% ± 6.8% 116,412	26.0% ± 5.5% 100,648	22.3% ± 4.4% 88,421	25.1% ± 5.6% 87,313
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.8% ± 2.7% 7,633	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.6% ± 1.7% 8,984

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	11.1% ± 10.5% 1,831	7.1% ± 8.0% 1,639	9.9% ± 11.7% 1,938	13.2% ± 13.0% 2,127	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	14.3% ± 6.7% 7,074	11.3% ± 5.9% 6,470	25.8% ± 9.0% 13,286	21.7% ± 8.6% 11,254	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Baltimore City	14.9% 1,435	14.6% 1,328	13.9% 1,254	13.4% 1,214	12.0% 1,106	11.6% 1,064	10.3% 1,009	9.9% 997	10.3% 1,025	8.8% 833

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	48.1% ± 6.9% 70,800	49.8% ± 6.5% 89,794	43.7% ± 6.2% 72,072	*32.7% ± 5.5% 58,066	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Baltimore County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	22.0% ± 4.1% 123,958	24.4% ± 4.6% 141,793	20.4% ± 3.6% 121,830	16.9% ± 3.1% 110,467	15.6% ± 3.8% 100,859
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	0.7% ± 0.8% 3,930	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.4% ± 1.8% 15,732

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	5.6% ± 6.2% 1,620	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	11.1% ± 12.8% 3,942	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	33.0% ± 9.8% 19,762	23.2% ± 10.5% 12,102	36.1% ± 13.7% 16,443	41.4% ± 11.4% 19,153	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Baltimore County	9.6% 904	8.9% 807	8.8% 791	8.8% 822	8.6% 809	7.8% 758	7.8% 776	7.6% 769	7.3% 748	6.9% 686

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	35.0% ± 5.4% 72,542	35.1% ± 5.4% 79,680	32.5% ± 6.2% 66,740	26.2% ± 4.8% 60,729	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Calvert County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	28.4% ± 10.8% 15,737	23.0% ± 8.0% 15,066	18.9% ± 6.1% 9,543	19.6% ± 6.9% 13,735	13.1% ± 6.5% 8,260
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.4% ± 4.2% 2,501	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.5% ± 3.0% 1,569

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	12.9% ± 12.6% 372	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	22.3% ± 11.6% 1,154	10.4% ± 10.1% 442	20.6% ± 13.9% 1,209	25.8% ± 13.2% 1,851	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Calvert County	14.2% 145	14.3% 137	12.7% 129	12.2% 129	11.8% 118	11.9% 119	12.7% 128	9.9% 97	11.5% 110	9.3% 86

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	38.9% ± 6.7% 10,387	41.8% ± 7.2% 10,673	30.5% ± 7.3% 8,805	30.7% ± 6.7% 10,497	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Caroline County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	24.8% ± 13.5% 7,390	DATA NOT AVAILABLE FOR 2002	25.6% ± 10.6% 6,257	19.3% ± 8.0% 4,709	29.9% ± 12.9% 8,495
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.7% ± 3.3% 505	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.2% ± 2.2% 627

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	24.5% ± 12.9% 461	17.5% ± 12.2% 265	NOT AVAILABLE— TOO FEW QUITTING RESPONDENTS	16.1% ± 14.7% 476	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Caroline County	17.0% 69	18.3% 71	15.1% 58	16.2% 68	15.9% 74	17.8% 76	12.8% 61	13.7% 64	12.8% 64	15.3% 68

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	43.2% ± 7.1% 4,237	34.9% ± 8.9% 3,103	31.9% ± 9.4% 3,109	32.3% ± 8.4% 3,433	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Carroll County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	14.5% ± 7.4% 15,323	22.6% ± 9.4% 26,267	12.3% ± 5.2% 19,376	12.0% ± 4.7% 14,424	21.2% ± 8.8% 33,754
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.4% ± 2.6% 1,432	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	4.5% ± 5.3% 7,239

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	24.7% ± 14.6% 2,092	50.0% ± 21.1% 3,776	29.6% ± 18.4% 2,438	36.2% ± 16.0% 2,768	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Carroll County	11.1% 211	10.6% 199	11.2% 213	9.6% 188	9.6% 191	9.6% 184	10.2% 191	11.1% 207	10.6% 186	9.9% 155

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	43.2% ± 7.1% 4,237	34.9% ± 8.9% 3,103	31.9% ± 9.4% 3,109	32.3% ± 8.4% 3,433	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Cecil County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	26.7% ± 8.0% 16,364	30.1% ± 9.4% 18,884	24.4% ± 6.0% 23,277	23.1% ± 8.6% 15,644	24.7% ± 7.3% 16,923
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	2.5% ± 3.6% 1,549	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.4% ± 2.2% 1,621

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	13.1% ± 10.5% 927	35.8% ± 14.9% 2,524	23.6% ± 10.3% 1,576	28.2% ± 12.8% 2,536	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Cecil County	23.3% 265	19.5% 230	18.2% 211	18.9% 230	19.7% 239	18.7% 227	19.7% 267	21.4% 283	22.8% 291	21.7% 257

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	40.5% ± 8.2% 10,313	45.1% ± 7.6% 13,240	39.7% ± 6.3% 12,514	37.2% ± 7.4% 12,995	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Charles County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	25.5% ± 7.1% 20,249	23.9% ± 7.9% 23,068	18.4% ± 6.2% 19,814	18.1% ± 6.9% 18,120	14.1% ± 4.7% 16,158
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	0.0% ± 0.0% 0	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	1.8% ± 2.2% 2,043

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	12.3% ± 11.1% 901	17.3% ± 13.5% 1,121	38.8% ± 15.0% 2,854	25.4% ± 11.2% 2,432	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Charles County	13.3% 232	12.2% 213	11.1% 195	9.4% 177	10.7% 194	7.1% 132	8.7% 169	6.8% 134	7.0% 134	6.6% 119

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	35.6% ± 7.9% 13,362	35.6% ± 7.2% 14,684	31.3% ± 6.1% 14,898	30.5% ± 5.6% 15,450	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Dorchester County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	26.4% ± 16.6% 6,452	33.7% ± 16.2% 10,384	21.0% ± 7.6% 5,005	20.1% ± 8.8% 4,485	13.1% ± 6.8% 2,885
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	2.6% ± 5.0% 635	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	1.8% ± 2.7% 392

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	20.3% ± 13.6% 669	12.8% ± 9.7% 271	<u>47.8%</u> ± 22.0% 983	27.6% ± 15.4% 801	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Dorchester County	18.6% 61	14.3% 48	16.6% 51	14.9% 52	13.2% 48	17.5% 57	13.1% 53	11.4% 47	14.4% 65	15.1% 62

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	35.6% ± 7.9% 13,362	35.6% ± 7.2% 14,684	31.3% ± 6.1% 14,898	30.5% ± 5.6% 15,450	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Frederick County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	20.6% ± 5.2% 29,019	18.7% ± 5.4% 26,100	18.7% ± 4.7% 31,574	13.8% ± 3.6% 22,906	12.1% ± 4.0% 19,405
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.6% ± 1.5% 2,211	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.6% ± 1.6% 4,122

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	19.2% ± 12.6% 2,326	24.7% ± 15.5% 2,660	25.7% ± 14.3% 1,908	24.5% ± 12.8% 3,007	DATA NOT AVAILABLE FOR 2010

1 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

2 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

3 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Frederick County	12.1% 351	12.0% 349	9.2% 276	9.6% 297	8.6% 254	8.0% 241	7.5% 231	7.3% 224	7.4% 219	8.4% 243

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	35.0% ± 7.0% 20,570	32.3% ± 7.1% 20,506	29.2% ± 5.6% 21,057	23.1% ± 5.2% 18,514	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Garrett County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	20.2% ± 13.7% 4,748	19.9% ± 11.8% 5,123	17.0% ± 6.3% 4,886	18.3% ± 6.7% 5,137	19.1% ± 9.1% 5,038
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	18.5% ± 13.4% 4,349	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	6.0% ± 5.1% 1,590

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	14.6% ± 12.1% 237	24.6% ± 15.7% 408	30.6% ± 21.6% 375	44.4% ± 21.8% 769	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Garrett County	16.8% 56	19.2% 63	17.0% 51	16.7% 57	20.9% 66	19.4% 61	16.2% 48	18.1% 57	17.0% 47	13.7% 42

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	23.3% ± 9.2% 2,250	36.5% ± 8.6% 2,955	30.8% ± 8.4% 2,709	21.2% ± 7.8% 1,809	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Harford County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	17.0% ± 6.8% 29,816	20.0% ± 7.5% 29,626	18.7% ± 5.9% 36,475	21.2% ± 6.7% 40,252	20.1% ± 6.8% 35,944
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	0.8% ± 1.6% 1,432	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	1.9% ± 2.1% 3,488

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	11.5% ± 13.6% 1,250	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	22.2% ± 15.9% 2,249	16.9% ± 15.2% 1,715	25.2% ± 10.3% 3,649	44.1% ± 19.6% 6,251	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Harford County	13.1% 386	10.1% 291	12.4% 360	11.6% 337	10.1% 301	10.7% 314	11.3% 342	10.8% 327	9.1% 271	8.4% 239

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	36.9% ± 7.3% 24,219	30.2% ± 6.8% 22,244	31.9% ± 5.1% 24,769	*20.6% ± 5.4% 16,855	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Howard County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	13.4% ± 5.6% 24,711	21.7% ± 8.3% 38,437	12.6% ± 4.4% 29,226	6.5% ± 3.4% 15,859	8.6% ± 4.7% 20,076
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	0.9% ± 1.7% 1,615	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	0.3% ± 0.6% 674

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	22.5% ± 12.3% 2,181	29.6% ± 15.6% 2,457	30.6% ± 15.3% 2,042	38.1% ± 15.0% 3,853	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Howard County	3.3% 119	4.4% 158	3.3% 117	3.6% 128	3.3% 116	3.7% 124	2.9% 99	2.7% 93	2.2% 74	1.4% 46

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	26.7% ± 5.3% 23,104	27.1% ± 5.4% 23,244	19.9% ± 4.4% 18,602	*15.3% ± 3.5% 14,026	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Kent County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	<u>19.4%</u> ± 7.8% 3,076	<u>15.1%</u> ± 6.2% 2,768	<u>13.4%</u> ± 6.8% 1,749
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	<u>1.0%</u> ± 1.9% 125

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	<u>32.2%</u> ± 20.1% 437	<u>23.4%</u> ± 17.2% 356	NOT AVAILABLE— TOO FEW QUITTING RESPONDENTS	<u>35.0%</u> ± 25.4% 598	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Kent County	21.2% 43	20.0% 31	16.6% 26	18.6% 31	16.3% 33	15.5% 28	19.0% 36	19.7% 35	13.7% 30	14.4% 27



ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	41.0% ± 9.2% 1,784	38.1% ± 9.9% 1,962	38.9% ± 11.8% 2,233	33.6% ± 11.7% 1,696	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶

(Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Montgomery County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	13.2% ± 3.1% 82,261	15.8% ± 4.1% 110,320	9.3% ± 2.4% 70,439	8.0% ± 2.8% 55,297	8.0% ± 2.6% 55,245
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.1% ± 0.9% 6,928	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	1.1% ± 0.7% 7,281

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	25.2% ± 13.0% 8,174	31.5% ± 17.3% 11,947	35.2% ± 16.5% 10,928	34.3% ± 13.6% 12,073	DATA NOT AVAILABLE FOR 2010

1 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

2 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

3 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates


POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Montgomery County	2.5% 327	2.0% 269	1.3% 168	1.3% 171	1.0% 142	0.9% 120	0.7% 95	0.5% 68	0.6% 80	0.6% 76

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	19.9% ± 4.4% 51,360	19.4% ± 4.7% 52,278	20.8% ± 4.5% 59,233	14.0% ± 3.4% 43,342	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Prince George's County Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	16.6% ± 3.7% 95,239	17.7% ± 4.3% 96,524	17.2% ± 3.8% 90,828	<u>11.5%</u> ± 2.8% 69,211	13.5% ± 3.5% 85,283
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	0.4% ± 0.6% 2,178	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	1.4% ± 1.4% 8,632

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	12.8% ± 13.5% 2,653	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	22.3% ± 12.3% 9,045	25.0% ± 11.7% 8,599	34.0% ± 13.8% 10,646	24.6% ± 10.4% 10,801	DATA NOT AVAILABLE FOR 2010

1 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

2 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

3 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Prince George's County	3.6% 447	3.1% 386	2.5% 316	2.1% 269	1.9% 237	1.6% 205	1.4% 177	1.4% 185	1.7% 213	1.7% 204

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	32.6% ± 5.7% 70,153	25.5% ± 5.7% 66,752	26.2% ± 4.5% 71,933	23.8% ± 4.5% 61,378	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



4 Source: Vital Statistics

5 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

6 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Queen Anne's County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	28.5% ± 13.0% 7,426	11.7% ± 8.4% 3,288	21.7% ± 6.4% 7,095	19.2% ± 6.7% 7,350	16.0% ± 6.4% 6,216
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	4.9% ± 6.7% 1,264	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	4.5% ± 4.3% 1,740

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	17.4% ± 10.4% 568	34.7% ± 16.9% 803	49.4% ± 28.9% 1,730	40.3% ± 17.4% 1,588	DATA NOT AVAILABLE FOR 2010

1 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

2 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

3 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Queen Anne's County	15.6% 78	11.7% 55	10.2% 54	8.0% 40	10.8% 55	9.8% 51	8.9% 46	11.3% 63	6.2% 33	10.0% 50

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	40.2% ± 7.9% 5,328	35.1% ± 7.5% 4,143	28.1% ± 8.6% 3,766	37.6% ± 9.1% 5,933	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Somerset County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	26.3% ± 10.3% 3,329	28.1% ± 11.4% 4,236	30.9% ± 15.9% 4,911
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	4.1% ± 4.3% 651

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	28.5% ± 13.8% 566	24.2% ± 14.1% 373	33.2% ± 30.2% 586	25.0% ± 18.7% 580	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Somerset County	20.9% 57	23.0% 56	17.0% 44	19.7% 51	16.5% 44	16.2% 43	16.1% 45	14.8% 37	14.8% 41	16.6% 41

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	44.8% ± 9.0% 3,226	37.4% ± 9.8% 2,268	46.6% ± 17.0% 3,055	37.3% ± 10.1% 2,268	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



St. Mary's County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	20.6% ± 7.9% 10,609	31.9% ± 10.0% 19,090	21.9% ± 7.3% 14,965	22.4% ± 6.7% 17,757	15.9% ± 5.2% 13,186
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	5.8% ± 6.0% 2,953	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	3.9% ± 3.8% 3,179

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	21.1% ± 11.3% 1,311	26.6% ± 17.0% 1,230	54.3% ± 17.2% 2,945	34.5% ± 17.5% 1,767	DATA NOT AVAILABLE FOR 2010

1 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

2 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

3 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
St. Mary's County	13.0% 158	14.3% 181	12.3% 165	11.8% 159	13.5% 195	11.5% 160	9.9% 148	10.5% 155	8.8% 127	10.0% 148

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	37.8% ± 6.9% 11,082	38.2% ± 7.4% 10,628	29.6% ± 7.1% 9,208	22.7% ± 5.6% 7,463	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Talbot County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	28.4% ± 15.3% 7,099	21.2% ± 11.8% 5,630	16.3% ± 5.7% 4,564	13.2% ± 5.3% 3,208	13.6% ± 6.3% 4,332
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	0.0% ± 0.0% 0	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	0.3% ± 0.5% 83

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	17.6% ± 12.8% 358	14.3% ± 11.8% 389	45.4% ± 23.3% 1,086	NOT AVAILABLE— TOO FEW QUITTING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Talbot County	14.4% 53	10.1% 36	10.2% 34	10.3% 35	8.0% 31	12.9% 47	10.1% 36	8.2% 30	10.6% 41	6.8% 24

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	31.1% ± 9.5% 2,476	41.3% ± 10.7% 3,210	31.8% ± 11.7% 3,107	29.7% ± 11.5% 2,980	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Washington County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	24.4% ± 6.5% 22,823	23.9% ± 6.2% 25,628	22.8% ± 5.2% 24,044	25.6% ± 6.2% 30,527	18.3% ± 5.2% 19,400
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	3.5% ± 2.9% 3,268	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	5.3% ± 3.9% 5,585

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW QUITTING RESPONDENTS	16.5% ± 15.2% 1,268	29.7% ± 13.6% 1,980	31.8% ± 16.3% 2,201	DATA NOT AVAILABLE FOR 2010

1 Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

2 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

3 Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk '*'.

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Washington County	19.1% 305	20.9% 339	18.1% 306	17.5% 296	15.6% 266	15.8% 278	16.3% 310	16.3% 322	16.3% 298	16.8% 294

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	40.5% ± 8.7% 15,369	31.2% ± 7.9% 12,602	28.7% ± 5.9% 12,548	25.3% ± 6.0% 11,071	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Wicomico County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	21.7% ± 8.2% 14,221	24.8% ± 8.4% 13,794	21.8% ± 6.7% 13,456	17.6% ± 6.3% 12,488	22.6% ± 7.6% 15,825
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	0.8% ± 1.6% 505	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	1.5% ± 1.4% 1,043

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	3.0% ± 3.8% 152	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	28.2% ± 14.6% 1,520	25.7% ± 14.0% 1,721	7.7% ± 7.8% 241	23.1% ± 15.7% 1,468	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Wicomico County	14.4% 169	16.6% 189	15.1% 175	15.7% 186	14.8% 172	16.7% 202	13.3% 177	15.2% 204	15.5% 212	13.6% 179

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	34.9% ± 9.4% 8,085	34.2% ± 8.1% 8,296	31.5% ± 8.6% 8,238	26.7% ± 7.0% 8,710	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.



Worcester County

Tobacco Use by Adults: 2000—2010

Maryland Department of Health and Mental Hygiene
Tobacco Use Prevention and Cessation Program

CURRENT USE OF TOBACCO PRODUCTS¹

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

TOBACCO PRODUCTS Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Any Tobacco Product (All Adult Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Any Tobacco Product (Adult Minority Populations)	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Cigarette	23.5% ± 10.1% 10,200	32.0% ± 11.4% 11,771	18.3% ± 7.6% 6,606	17.6% ± 9.6% 6,797	11.7% ± 7.2% 4,488
Cigar	DATA NOT AVAILABLE FOR 2000	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	DATA NOT AVAILABLE FOR 2010
Smokeless Tobacco	1.5% ± 2.8% 635	DATA NOT AVAILABLE FOR 2002	DATA NOT AVAILABLE FOR 2006	DATA NOT AVAILABLE FOR 2008	2.8% ± 2.8% 1,085

INITIATION OF TOBACCO USE IN PAST YEAR²

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

INITIATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	NOT AVAILABLE— TOO FEW INITIATING RESPONDENTS	DATA NOT AVAILABLE FOR 2010			

CESSATION OF TOBACCO USE IN PAST YEAR³

Current Adult (Age 18 and Older) Use of Tobacco Products
Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

CESSATION Prevalence (%), CI, and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Adult Population	13.9% ± 11.3% 339	10.4% ± 8.6% 361	36.4% ± 20.6% 1,158	NOT AVAILABLE— TOO FEW QUITTING RESPONDENTS	DATA NOT AVAILABLE FOR 2010

¹ Source: Behavioral Risk Factor Surveillance Survey, 2000, 2002, 2006, 2008, and 2010. The Behavioral Risk Factor Surveillance Survey collects data on use of cigarettes and smokeless tobacco only. It does not collect data on use of cigars or any other tobacco product. Thus, estimates of these behaviors are unavailable for 2000, 2002, 2006, 2008, and 2010. Estimates for use of smokeless tobacco are available for 2000 and 2010 only.

² Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

³ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined, between 2000 and 2010 preceded by an asterisk (**).

CURRENT CIGARETTE SMOKING — Pregnant Women⁴

Proportion (%) of Selected Populations and Number for Pregnant Women from 100% of Birth Certificates



POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2001	2002	2003	2004	2005	2006	2007	2008	2009
Maryland Statewide	9.2% 8,842	8.7% 8,361	8.0% 7,879	7.7% 7,759	7.4% 7,508	6.9% 7,001	6.8% 7,267	6.6% 7,152	6.6% 7,113	6.1% 6,593
Worcester County	16.9% 84	13.8% 72	15.4% 70	16.4% 77	14.9% 67	12.1% 60	16.1% 75	14.0% 70	14.2% 68	10.7% 49

ADULT HOUSEHOLDS WITH MINOR CHILDREN⁵

Current Adult (Age 18 and Older), Proportion (%), Confidence Interval (CI), and Estimated Number of Adults

POPULATIONS Prevalence (%) and Number Adults	2000 Baseline Data	2002	2006	2008	2010
Proportion with Adult Smokers	34.3% ± 16.6% 4,627	44.8% ± 10.0% 4,366	37.5% ± 11.1% 4,974	42.9% ± 13.1% 5,445	DATA NOT AVAILABLE FOR 2010

MOST IMPORTANT REASON FOR WANTING TO QUIT SMOKING CIGARETTES, 2008⁶ (Top 3 Reported as 'Most Important Reason' by Former Smokers and Smokers Trying to Quit)



⁴ Source: Vital Statistics

⁵ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.

⁶ Source: Maryland Adult Tobacco Survey, 2000, 2002, 2006, and 2008.